

Status: Currently Official on 16-Feb-2025
 Official Date: Official as of 01-Oct-2017
 Document Type: Reagents
 DocId: GUID-FB996016-022F-4955-A397-6A77A78FD23C_5_en-US
 DOI: https://doi.org/10.31003/USPNF_R3401_05_01
 DOI Ref: etm0o

© 2025 USPC
 Do not distribute

1 N Sulfuric Acid VS

H_2SO_4 , 98.08
 49.04 g in 1000 mL

Add slowly, with stirring, 27 mL of [sulfuric acid](#) to a sufficient quantity of [water](#) to make 1000 mL.

Standardization: Accurately weigh about 5.0 g of [tromethamine](#), dried according to the label instructions or, if this information is not available, dried at 105° for 3 h. Dissolve in 50 mL of [water](#) and add 2 drops of [bromocresol green TS](#). Titrate with 1 N sulfuric acid to a pale yellow endpoint. Each 121.14 mg of tromethamine is equivalent to 1 mL of 1 N sulfuric acid.

$$N = \frac{\text{mg tromethamine}}{121.14 \times \text{mL } H_2SO_4}$$

[NOTE—If this volumetric solution is used in a qualitative application such as pH adjustment, dissolution medium, or diluent, its standardization is not required.]

Auxiliary Information - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
1 N SULFURIC ACID VS	Margareth R.C. Marques Principal Scientific Liaison	HDQ Headquarters

Most Recently Appeared In:

Pharmacopeial Forum: Volume No. PF 42(1)

Current DocID: [GUID-FB996016-022F-4955-A397-6A77A78FD23C_5_en-US](#)

Previous DocID: [GUID-FB996016-022F-4955-A397-6A77A78FD23C_2_en-US](#)

Previous DocID: [GUID-FB996016-022F-4955-A397-6A77A78FD23C_4_en-US](#)

DOI: https://doi.org/10.31003/USPNF_R3401_05_01

DOI ref: [etm0o](#)