

Status: Currently Official on 12-Feb-2025
Official Date: Official as of 01-Dec-2017
Document Type: General Chapter
DocId: GUID-A160D6F1-944D-4136-83AE-8DD835CDB2E9_1_en-US
DOI: https://doi.org/10.31003/USPNF_M98865_01_01
DOI Ref: 3n6hx

© 2025 USPC
Do not distribute

⟨115⟩ DEXPANTHENOL ASSAY

INTRODUCTION

The following liquid chromatographic procedure is provided for the determination of dexpanthenol as a component in dietary supplements or pharmaceutical dosage forms.

While conducting this procedure, protect solutions containing and derived from the test specimen and the Reference Standards from the atmosphere and light, preferably with the use of low-actinic glassware. Use the appropriate USP Reference Standards.

ASSAY

• CHROMATOGRAPHIC METHODS, PROCEDURE 1

This procedure can be used to determine dexpanthenol in:

- [Oil- and Water-Soluble Vitamins Oral Solution](#)
- [Oil- and Water-Soluble Vitamins with Minerals Oral Solution](#)
- [Water-Soluble Vitamins with Minerals Oral Solution](#)

Unless specified in the monograph, the *Standard solution*, *System suitability solution*, and *Sample solution* are prepared as follows.

Mobile phase: 0.2 M monobasic sodium phosphate and [methanol](#) (97:3). Adjust with 1.7 M [phosphoric acid](#) to a pH of 3.2 ± 0.1 .

Standard solution: 80 µg/mL of [USP Dexpanthenol RS](#) in *Mobile phase*

System suitability solution: 80 µg/mL of [USP Calcium Pantothenate RS](#) in *Mobile phase*. Mix the resulting solution and *Standard solution* (1:1).

Sample solution: Equivalent to 80 µg/mL of dexpanthenol from Oral Solution in *Mobile phase*

Chromatographic system

(See [Chromatography \(621\)](#), [System Suitability](#).)

Mode: LC

Detector: UV 210 nm

Column: 4.0-mm × 10-cm; packing [L1](#)

Flow rate: 1 mL/min

Injection volume: 20 µL

System suitability

Samples: *Standard solution* and *System suitability solution*

Suitability requirements

Resolution: NLT 1.5 between dexpanthenol and calcium pantothenate, *System suitability solution*

Tailing factor: NMT 2.0 for the dexpanthenol peak, *Standard solution*

Relative standard deviation: NMT 2.0%, *Standard solution*

Analysis

Samples: *Standard solution* and *Sample solution*

Calculate the percentage of the labeled amount of dexpanthenol ($C_9H_{19}NO_4$) in the portion of the sample taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times 100$$

r_U = peak response of dexpanthenol from the *Sample solution*

r_S = peak response of dexpanthenol from the *Standard solution*

C_S = concentration of [USP Dexpanthenol RS](#) in the *Standard solution* (mg/mL)

C_U = nominal concentration of dexpanthenol in the *Sample solution* (mg/mL)

ADDITIONAL REQUIREMENTS

• [USP Reference Standards \(11\)](#)

[USP Calcium Pantothenate RS](#)

[USP Dexpanthenol RS](#)

Auxiliary Information - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
<115> DEXPANTHENOL ASSAY	Natalia Davydova Scientific Liaison	NBDS2020 Non-botanical Dietary Supplements

Most Recently Appeared In:

Pharmacopeial Forum: Volume No. PF 42(4)

Current DocID: GUID-A160D6F1-944D-4136-83AE-8DD835CDB2E9_1_en-US

DOI: https://doi.org/10.31003/USPNF_M98865_01_01

DOI ref: [3n6hx](#)

OFFICIAL