

Status: Currently Official on 17-Feb-2025
Official Date: Official Prior to 2013
Document Type: Reagents
DocId: GUID-81C62626-BF7C-460C-B804-7C13FC607942_1_en-US
DOI: https://doi.org/10.31003/USPNF_R1579_01_01
DOI Ref: 4e4ea

© 2025 USPC
Do not distribute

2,4,6-Triamino-5-nitrosopyrimidine,

$C_4H_6N_6O$ 154.13—Pink powder.

Assay: Dissolve about 34 mg, accurately weighed, in 50 mL of glacial acetic acid. Titrate with 0.1 N perchloric acid VS, determining the endpoint potentiometrically. Perform a blank determination, and make any necessary corrections. Each mL of 0.1 N perchloric acid is equivalent to 15.41 mg of $C_4H_6N_6O$. Not less than 97% is found.

Auxiliary Information - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
2,4,6-TRIAMINO-5-NITROSOPYRIMIDINE	Margareth R.C. Marques Principal Scientific Liaison	HDQ Headquarters

Most Recently Appeared In:

Pharmacopeial Forum: Volume No. Information currently unavailable

Current DocID: [GUID-81C62626-BF7C-460C-B804-7C13FC607942_1_en-US](#)

DOI: https://doi.org/10.31003/USPNF_R1579_01_01

DOI ref: [4e4ea](#)

OFFICIAL