

Status: Currently Official on 14-Feb-2025
Official Date: Official Prior to 2013
Document Type: Reagents
DocId: GUID-B38E5DAC-D5F1-473E-90BD-429E8A26B0DB_1_en-US
DOI: https://doi.org/10.31003/USPNF_R3155_01_01
DOI Ref: 66j68

© 2025 USPC
Do not distribute

2,7-Dihydroxynaphthalene TS

—Dissolve 100 mg of 2,7-dihydroxynaphthalene in 1000 mL of sulfuric acid, and allow the solution to stand until the yellow color disappears. If the solution is very dark, discard it and prepare a new solution from a different supply of sulfuric acid. This solution is stable for approximately 1 month if stored in a dark bottle.

Auxiliary Information - Please [check for your question in the FAQs](#) before contacting USP.

| Topic/Question | Contact | Expert Committee |
|-----------------------------|--|------------------|
| 2,7-DIHYDROXYNAPHTHALENE TS | Margareth R.C. Marques Principal Scientific Liaison | HDQ Headquarters |

Most Recently Appeared In:

Pharmacopeial Forum: Volume No. Information currently unavailable

Current DocId: [GUID-B38E5DAC-D5F1-473E-90BD-429E8A26B0DB_1_en-US](#)

DOI: https://doi.org/10.31003/USPNF_R3155_01_01

DOI ref: [66j68](#)

OFFICIAL