

Status: Currently Official on 16-Feb-2025  
Official Date: Official as of 01-Nov-2020  
Document Type: Reagents  
DocId: GUID-443EC259-A10C-4492-A965-0D5525AFED55\_2\_en-US  
DOI: [https://doi.org/10.31003/USPNF\\_R8022\\_02\\_01](https://doi.org/10.31003/USPNF_R8022_02_01)  
DOI Ref: tq7x8

© 2025 USPC  
Do not distribute

**Add the following:**

## ^50% Sodium Hydroxide TS

—Transfer 500 g of [sodium hydroxide](#) to a 1000-mL volumetric flask. Dissolve in and dilute with [carbon dioxide-free water](#) to volume. Allow to stand. Decant the clear supernatant, taking precautions to avoid the introduction of carbon dioxide.▲ (USP 1-May-2020)

**Auxiliary Information** - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
50 PERCENT SODIUM HYDROXIDE TS	<a href="#">Margareth R.C. Marques</a> Principal Scientific Liaison	HDQ Headquarters

**Most Recently Appeared In:**

Pharmacopeial Forum: Volume No. PF 44(6)

**Current DocID:** [GUID-443EC259-A10C-4492-A965-0D5525AFED55\\_2\\_en-US](#)

**DOI:** [https://doi.org/10.31003/USPNF\\_R8022\\_02\\_01](https://doi.org/10.31003/USPNF_R8022_02_01)

**DOI ref:** [tq7x8](#)

OFFICIAL