

Status: Currently Official on 17-Feb-2025
Official Date: Official as of 01-Dec-2016
Document Type: USP Monographs
DocId: GUID-6E40BCFA-5DD2-46BC-AFE6-99A2B1F34A33_1_en-US
DOI: https://doi.org/10.31003/USPNF_M1376_01_01
DOI Ref: u452u

© 2025 USPC
Do not distribute

Acetazolamide Compounded Oral Suspension

DEFINITION

Acetazolamide Compounded Oral Suspension contains NLT 90.0% and NMT 110.0% of the labeled amount of acetazolamide ($C_4H_6N_4O_3S_2$).
Prepare Acetazolamide Compounded Oral Suspension, 25 mg/mL, as follows (see [Pharmaceutical Compounding—Nonsterile Preparations \(795\)](#)).

Acetazolamide	2.5 g
Vehicle: a 1:1 mixture of Vehicle for Oral Solution, <i>NF</i> (regular or sugar-free), and Vehicle for Oral Suspension, <i>NF</i> , or Cherry Syrup, <i>NF</i> , a sufficient quantity to make	100 mL

If using tablets, place in a mortar and comminute to a fine powder, or add *Acetazolamide* powder. Add about 20 mL of the *Vehicle*, and mix to a uniform paste. Add the *Vehicle* in small portions almost to volume, and mix thoroughly after each addition. Transfer the contents of the mortar, stepwise and quantitatively, to a calibrated bottle. Add enough liquid *Vehicle* to bring to final volume, and mix well.

ASSAY

PROCEDURE

Mobile phase: Dissolve 4.1 g of anhydrous sodium acetate in 950 mL of water, and add 20 mL of methanol and 30 mL of acetonitrile. Adjust with glacial acetic acid to a pH of 4.0.

Standard stock solution: Transfer about 25 mg of [USP Acetazolamide RS](#), accurately weighed, to a 50-mL volumetric flask, add 5.0 mL of 0.5 N sodium hydroxide, and mix to dissolve. Dilute with water to volume, and mix.

Standard solution: 250 µg/mL of [USP Acetazolamide RS](#) from the *Standard stock solution* in water

Sample solution: 250 µg/mL of acetazolamide from Oral Suspension in *Mobile phase*. Agitate the container of Oral Suspension for 30 min on a rotating mixer, remove a 5-mL sample, and store in a clear glass vial at −70° until analyzed. At the time of analysis, remove the sample from the freezer, allow to reach room temperature, and mix with a vortex mixer for 30 s. Pipet 1.0 mL of this solution to a 100-mL volumetric flask, and dilute with *Mobile phase* to volume.

Chromatographic system

(See [Chromatography \(621\), System Suitability](#).)

Mode: LC

Detector: UV 254 nm

Column: 4.6-mm × 25-cm; 5-µm packing L1

Flow rate: 2 mL/min

Injection volume: 20 µL

System suitability

Sample: *Standard solution*

[NOTE—The retention time for the acetazolamide peak is about 3 min.]

Suitability requirements

Relative standard deviation: NMT 1.1% for replicate injections

Analysis

Samples: *Standard solution* and *Sample solution*

Calculate the percentage of the labeled amount of acetazolamide ($C_4H_6N_4O_3S_2$) in the portion of Oral Suspension taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times 100$$

r_U = peak response from the *Sample solution*

r_S = peak response from the *Standard solution*

C_S = concentration of [USP Acetazolamide RS](#) in the *Standard solution* (µg/mL)

C_U = nominal concentration of acetazolamide in the *Sample solution* (µg/mL)

Acceptance criteria: 90.0%–110.0%

SPECIFIC TESTS

- **pH (791):** 4.0–5.0 (Vehicle for Oral Solution and Vehicle for Oral Suspension), 3.1–3.9 (Cherry Syrup)

ADDITIONAL REQUIREMENTS

- **PACKAGING AND STORAGE:** Package in tight, light-resistant containers. Store at controlled room temperature, or in a refrigerator.
- **BEYOND-USE DATE:** NMT 60 days after the day on which it was compounded when stored at controlled room temperature, or in a refrigerator
- **LABELING:** Label it to state that it is to be well shaken before use, and to state the *Beyond-Use Date*.
- **USP REFERENCE STANDARDS (11).**
[USP Acetazolamide RS](#)

Auxiliary Information - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
ACETAZOLAMIDE COMPOUNDED ORAL SUSPENSION	Brian Serumaga Science Program Manager	CMP2020 Compounding 2020
REFERENCE STANDARD SUPPORT	RS Technical Services RSTECH@usp.org	CMP2020 Compounding 2020

Chromatographic Database Information: [Chromatographic Database](#)

Most Recently Appeared In:

Pharmacopeial Forum: Volume No. PF 40(5)

Current DocID: GUID-6E40BCFA-5DD2-46BC-AFE6-99A2B1F34A33_1_en-US

DOI: https://doi.org/10.31003/USPNF_M1376_01_01

DOI ref: [u452u](#)

OFFICIAL