

Status: Currently Official on 17-Feb-2025
Official Date: Official as of 01-May-2020
Document Type: USP Monographs
DocId: GUID-533B5F61-CDDD-42CB-9AA3-411AB1C0757E_4_en-US
DOI: https://doi.org/10.31003/USPNF_M2950_04_01
DOI Ref: g5mcp

© 2025 USPC
Do not distribute

Aminocaproic Acid Tablets

DEFINITION

Aminocaproic Acid Tablets contain NLT 95.0% and NMT 105.0% of the labeled amount of aminocaproic acid ($C_6H_{13}NO_2$).

IDENTIFICATION

Change to read:

- **A.** ▲ [SPECTROSCOPIC IDENTIFICATION TESTS \(197\)](#), [Infrared Spectroscopy: 197K](#)▲ (CN 1-MAY-2020)

Sample: Triturate 2 Tablets with 10 mL of [water](#), and filter into 100 mL of [acetone](#). Swirl the mixture, and allow to stand for 15 min to complete crystallization. Pass the solution through a sintered-glass filter of medium pore size, and wash the crystals with 25 mL of [acetone](#). Apply vacuum to remove the solvent, dry at 105° for 30 min, and cool. Use the residue.

Acceptance criteria: Meet the requirements

ASSAY

• PROCEDURE

Sample solution: Nominally equivalent to about 500 mg of aminocaproic acid from NLT 20 finely powdered Tablets taken in a beaker in about 100 mL of [glacial acetic acid](#). Heat gently to effect solution, and cool.

Titrimetric system

Mode: Direct titration

Titrant: [0.1 N perchloric acid in dioxane VS](#)

Endpoint detection: Visual

Analysis: To the *Sample solution* add 10 drops of a 1-in-500 solution of [crystal violet](#) in [chlorobenzene](#). Titrate with *Titrant* to a blue endpoint, and perform a blank determination. Each mL of 0.1 N perchloric acid is equivalent to 13.12 mg of aminocaproic acid ($C_6H_{13}NO_2$).

Acceptance criteria: 95.0%–105.0%

PERFORMANCE TESTS

• [DISSOLUTION \(711\)](#)

Test 1

Medium: [Water](#); 900 mL

Apparatus 1: 100 rpm

Time: 45 min

Buffer: Dissolve 6.185 g of [boric acid](#) and 7.930 g of [potassium chloride](#) in about 1000 mL of [water](#), and add 60 mL of 1.0 N sodium hydroxide. Dilute with [water](#) to 2000 mL, and adjust if necessary with 1.0 N sodium hydroxide to a pH of 9.5 ± 0.1 .

Standard solution: 0.5 mg/mL of [USP Aminocaproic Acid RS](#) in [water](#)

Sample solution: Filter a portion of the solution under test.

Blank: [Water](#)

Analysis: Into three separate 50-mL volumetric flasks pipet 1 mL each of *Sample solution*, *Standard solution*, and *Blank*. Add 20.0 mL of *Buffer* and 3.0 mL of a freshly prepared 1-in-500 solution of [β-naphthoquinone-4-sodium sulfonate](#) to each flask. Swirl to mix, and place the three flasks in a water bath maintained at a temperature of $65 \pm 5^\circ$ for 45 min. Cool, and dilute the contents of each flask with [water](#) to volume.

Determine the percentage of the labeled amount of aminocaproic acid ($C_6H_{13}NO_2$) dissolved from absorbances, at the wavelength of maximum absorbance at about 460 nm, from the *Sample solution* in comparison with those from the *Standard solution*, using the *Blank* to set the instrument.

Tolerances: NLT 75% (Q) of the labeled amount of aminocaproic acid ($C_6H_{13}NO_2$) is dissolved.

Test 2: If the product complies with this test, the labeling indicates that it meets USP *Dissolution Test 2*.

Medium: 0.1 N hydrochloric acid; 500 mL

Apparatus 1: 100 rpm

Time: 30 min

Buffer A: Dissolve 500 mg of [sodium 1-heptanesulfonate](#) in 1 L of [water](#). Add 1.0 mL of [triethylamine](#) and mix well.

Buffer B: Dissolve 13.3 g of [monobasic sodium phosphate](#) in 1 L of *Buffer A*, and mix well. Adjust with [phosphoric acid](#) to a pH of 2.20 ± 0.05 .

[NOTE—The pH of *Buffer B* is critical because the diluent peak can coelute with the main peak even when the pH of *Buffer B* is at 2.10 or 2.30.]

Mobile phase: [Methanol](#) and *Buffer B* (25:75)

Standard solution: 1 mg/mL of [USP Aminocaproic Acid RS](#) in *Medium*. Sonication may be needed to aid the dissolution.

Sample solution: Pass a portion of the solution under test through a suitable filter of 0.45-µm pore size. Discard the first few milliliters of filtrate.

Chromatographic system

(See [Chromatography \(621\)](#), [System Suitability](#).)

Mode: LC

Detector: UV 210 nm

Column: 4.6 mm × 25 cm; 5-µm packing [L1](#)

Temperatures

Autosampler: 25°

Column: 50°

Flow rate: 1 mL/min

Injection volume: 25 µL

Run time: NLT 2.5 times the retention time of aminocaproic acid

System suitability

Sample: *Standard solution*

Suitability requirements

Tailing factor: NMT 2.5

Relative standard deviation: NMT 2.0%

Analysis

Samples: *Standard solution* and *Sample solution*

Calculate the percentage of the labeled amount of aminocaproic acid (C₆H₁₃NO₂) dissolved:

$$\text{Result} = (r_U/r_S) \times C_S \times V \times (1/L) \times 100$$

r_U = peak response of aminocaproic acid from the *Sample solution*

r_S = peak response of aminocaproic acid from the *Standard solution*

C_S = concentration of [USP Aminocaproic Acid RS](#) in the *Standard solution* (mg/mL)

V = volume of *Medium*, 500 mL

L = label claim (mg/Tablet)

Tolerances: NLT 80% (Q) of the labeled amount of aminocaproic acid (C₆H₁₃NO₂) is dissolved.

- [UNIFORMITY OF DOSAGE UNITS \(905\)](#): Meet the requirements

ADDITIONAL REQUIREMENTS

- **PACKAGING AND STORAGE:** Preserve in tight containers.
- **LABELING:** When more than one *Dissolution* test is given, the labeling states the *Dissolution* test used only if *Test 1* is not used.
- [USP REFERENCE STANDARDS \(11\)](#),
[USP Aminocaproic Acid RS](#)

Auxiliary Information - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
AMINOCAPROIC ACID TABLETS	Documentary Standards Support	SM22020 Small Molecules 2
REFERENCE STANDARD SUPPORT	RS Technical Services RSTECH@usp.org	SM22020 Small Molecules 2

Chromatographic Database Information: [Chromatographic Database](#)

Most Recently Appeared In:

Pharmacopeial Forum: Volume No. 50(6)

Current DocID: [GUID-533B5F61-CDDD-42CB-9AA3-411AB1C0757E_4_en-US](#)

DOI: https://doi.org/10.31003/USPNF_M2950_04_01

OFFICIAL