

Status: Currently Official on 17-Feb-2025
Official Date: Official as of 01-May-2024
Document Type: USP Monographs
DocId: GUID-26183518-D649-435D-8C26-045F7CD2B9FB_10_en-US
DOI: https://doi.org/10.31003/USPNF_M1069_10_01
DOI Ref: uiyfk

© 2025 USPC
Do not distribute

Amlodipine and Atorvastatin Tablets

DEFINITION

Amlodipine and Atorvastatin Tablets contain an amount of amlodipine besylate equivalent to NLT 90.0% and NMT 110.0% of the labeled amount of amlodipine ($C_{20}H_{25}ClN_2O_5$) and an amount of atorvastatin calcium equivalent to NLT 94.5% and NMT 105.0% of the labeled amount of atorvastatin ($C_{33}H_{35}FN_2O_5$). It may contain suitable antioxidants.

IDENTIFICATION

- **A.** The UV spectrum of the major peaks of the *Sample solution* exhibits maxima and minima at the same wavelengths as that of the *Standard solution*, as obtained in the Assay.
- **B.** The retention times of the major peaks of the *Sample solution* correspond to those of the *Standard solution*, as obtained in the Assay.

ASSAY

Change to read:

• **PROCEDURE**

Solution A: Dissolve 1.54 g of [ammonium acetate](#) in 1000 mL of [water](#) and add 2 mL of [triethylamine](#). Adjust with [acetic acid](#) to a pH of 5.0.

Mobile phase: [Acetonitrile](#), [methanol](#), and *Solution A* (38:15:47)

Buffer: Transfer 7 mL of [triethylamine](#) to a 1000-mL volumetric flask containing 900 mL of [water](#) and mix. Adjust with dilute [phosphoric acid](#) (1 in 100) to a pH of 3.0 and dilute with [water](#) to volume.

Diluent: [Acetonitrile](#), [methanol](#), and *Buffer* (3:7:10)

Standard stock solution 1: 0.35 mg/mL of [USP Amlodipine Besylate RS](#) in [methanol](#)

Standard stock solution 2: 0.44 mg/mL of [USP Atorvastatin Calcium RS](#) in [methanol](#)

Standard solution: Prepare solutions of [USP Amlodipine Besylate RS](#) and [USP Atorvastatin Calcium RS](#) in *Mobile phase* at concentrations given in [Table 1](#) from *Standard stock solution 1* and *Standard stock solution 2*.

Table 1

Strength of Tablet, Amlodipine/Atorvastatin (mg/mg)	Concentration of Amlodipine Besylate (mg/mL)	Concentration of Atorvastatin Calcium (mg/mL)
2.5/10, 5/20, 10/40	0.028	0.088
2.5/20, 5/40, 10/80	0.014	0.088
5/10, 10/20	0.028	0.044
2.5/40, 5/80	0.014	0.176
10/10	0.028	0.022

Sample solution: Transfer NLT 10 Tablets to a suitable volumetric flask. Add about 20% of the final volume of the volumetric flask size in *Diluent* and sonicate to disperse the Tablets. Add about 40% of the final volume of the volumetric flask size in *Diluent*, sonicate for 20 min, and dilute with *Diluent* to volume. Centrifuge and transfer a suitable quantity of the supernatant to an appropriate suitable volumetric flask. Dilute with *Mobile phase* to volume to obtain the nominal concentrations of amlodipine and atorvastatin similar to that of the *Standard solution*.

Chromatographic system

(See [Chromatography \(621\), System Suitability](#).)

Mode: LC

Detector: UV 237 nm. For *Identification A*, use a diode array detector in the range of 200–400 nm.

Column: 4.6-mm × 15-cm; 5-μm packing [L1](#)

Column temperature: 35°

Flow rate: 1 mL/min

Injection volume: 20 µL**Run time:** NLT 3.5 times the retention time of amlodipine**System suitability****Sample:** *Standard solution***Suitability requirements****Tailing factor:** NMT 2.0 for both peaks**Relative standard deviation:** NMT 2.0% for both peaks**Analysis****Samples:** *Standard solution* and *Sample solution*Calculate the percentage of the labeled amount of amlodipine ($C_{20}H_{25}ClN_2O_5$) in the portion of Tablets taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times (M_{r1}/M_{r2}) \times 100$$

 r_U = peak response of amlodipine from the *Sample solution* r_S = peak response of amlodipine from the *Standard solution* C_S = concentration of [USP Amlodipine Besylate RS](#) in the *Standard solution* (mg/mL) C_U = nominal concentration of amlodipine in the *Sample solution* (mg/mL) M_{r1} = molecular weight of amlodipine, 408.88 M_{r2} = molecular weight of amlodipine besylate, 567.05Calculate the percentage of the labeled amount of atorvastatin ($C_{33}H_{35}FN_2O_5$) in the portion of Tablets taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times [M \times (M_{r1}/M_{r2})] \times 100$$

 r_U = peak response of atorvastatin from the *Sample solution* r_S = peak response of atorvastatin from the *Standard solution* C_S = concentration of [USP Atorvastatin Calcium RS](#) in the *Standard solution* (mg/mL) C_U = nominal concentration of atorvastatin in the *Sample solution* (mg/mL) M = number of moles of atorvastatin per mole of atorvastatin calcium, 2 M_{r1} = molecular weight of atorvastatin, ▲558.65▲ (CN 1-May-2024) M_{r2} = molecular weight of atorvastatin calcium, ▲1155.36▲ (CN 1-May-2024)**Acceptance criteria****Amlodipine:** 90.0%–110.0%**Atorvastatin:** 94.5%–105.0%**PERFORMANCE TESTS****Change to read:**• [DISSOLUTION <711>](#)**Test 1****Solution A, Mobile phase, Standard stock solution 1, Standard stock solution 2, Chromatographic system, and System suitability:** Proceed as directed in the Assay.**Medium:** 0.1% [polysorbate 80](#) in [pH 6.8 phosphate buffer](#); 900 mL**Apparatus 2:** 75 rpm**Time:** 20 min**Standard solution:** ($L_1/900$) mg/mL of amlodipine and ($L_2/900$) mg/mL of atorvastatin in *Medium* from *Standard stock solution 1* and*Standard stock solution 2*, where L_1 is the label claim of amlodipine in mg/Tablet and L_2 is the label claim of atorvastatin in mg/Tablet**Sample solution:** Centrifuge the solution under test and use the supernatant.**Analysis****Samples:** *Standard solution* and *Sample solution*Calculate the percentage of the labeled amount of amlodipine ($C_{20}H_{25}ClN_2O_5$) dissolved:

$$\text{Result} = (r_U/r_S) \times C_S \times V \times (M_{r1}/M_{r2}) \times (1/L) \times 100$$

r_U = peak response of amlodipine from the *Sample solution*

r_S = peak response of amlodipine from the *Standard solution*

C_S = concentration of [USP Amlodipine Besylate RS](#) in the *Standard solution* (mg/mL)

V = volume of *Medium*, 900 mL

M_{r1} = molecular weight of amlodipine, 408.88

M_{r2} = molecular weight of amlodipine besylate, 567.05

L = label claim of amlodipine (mg/Tablet)

Calculate the percentage of the labeled amount of atorvastatin ($C_{33}H_{35}FN_2O_5$) dissolved:

$$\text{Result} = (r_U/r_S) \times C_S \times V \times [M \times (M_{r1}/M_{r2})] \times (1/L) \times 100$$

r_U = peak response of atorvastatin from the *Sample solution*

r_S = peak response of atorvastatin from the *Standard solution*

C_S = concentration of [USP Atorvastatin Calcium RS](#) in the *Standard solution* (mg/mL)

V = volume of *Medium*, 900 mL

M = number of moles of atorvastatin per mole of atorvastatin calcium, 2

M_{r1} = molecular weight of atorvastatin, 558.65▲ (CN 1-May-2024)

M_{r2} = molecular weight of atorvastatin calcium, 1155.36▲ (CN 1-May-2024)

L = label claim of atorvastatin (mg/Tablet)

Tolerances: NLT 80% (Q) of the labeled amount of amlodipine ($C_{20}H_{25}ClN_2O_5$) and atorvastatin ($C_{33}H_{35}FN_2O_5$) are dissolved.

Test 2: If the product complies with this test, the labeling indicates that it meets USP *Dissolution Test 2*.

Medium: 0.05 M phosphate buffer prepared as follows. Dissolve 40.8 g of [potassium phosphate, monobasic](#) and 5 g of [sodium hydroxide](#) in 6 L of [water](#). Adjust with [1 N sodium hydroxide](#) or [phosphoric acid](#) to a pH of 6.8; 900 mL.

Apparatus 2: 75 rpm

Time: 30 min

Buffer: Dissolve 4.0 g of [sodium phosphate, monobasic](#) in 1200 mL of [water](#) and add 6 mL of [triethylamine](#). Adjust with [phosphoric acid](#) to a pH of 2.5.

Mobile phase: [Acetonitrile](#) and *Buffer* (40:60)

Diluent: [Acetonitrile](#) and [water](#) (50:50)

Standard stock solution 1: 155 µg/mL of [USP Amlodipine Besylate RS](#) in *Diluent*

Standard stock solution 2: 460 µg/mL of [USP Atorvastatin Calcium RS](#) in *Diluent*

Standard solution: Prepare solutions of [USP Amlodipine Besylate RS](#) and [USP Atorvastatin Calcium RS](#) at concentrations given in [Table 2](#) from *Standard stock solution 1* and *Standard stock solution 2*.

Table 2

Strength of Tablet, Amlodipine/Atorvastatin (mg/mg)	Volume of Standard Stock Solution 1 to Be Added (mL)	Volume of Standard Stock Solution 2 to Be Added (mL)	Volume of Diluent to Be Added (mL)	Final Volume with Medium (mL)	Final Concentration of USP Amlodipine Besylate RS/ USP Atorvastatin Calcium RS (µg/mL)
5/10	10	5	45	200	7.75/11.5
5/20	10	10	40	200	7.75/23
5/40	10	20	30	200	7.75/46
5/80	10	40	10	200	7.75/92
10/10	20	5	35	200	15.5/11.5
10/20	20	10	30	200	15.5/23

Strength of Tablet, Amlodipine/Atorvastatin (mg/mg)	Volume of Standard Stock Solution 1 to Be Added (mL)	Volume of Standard Stock Solution 2 to Be Added (mL)	Volume of Diluent to Be Added (mL)	Final Volume with Medium (mL)	Final Concentration of USP Amlodipine Besylate RS/ USP Atorvastatin Calcium RS (µg/mL)
10/40	20	20	20	200	15.5/46
10/80	20	40	–	200	15.5/92

Sample solution: Pass a portion of the solution under test through a suitable filter. Discard the first few milliliters of filtrate.

Chromatographic system

(See [Chromatography \(621\), System Suitability](#).)

Mode: LC

Detector: UV 240 nm

Column: 4.6-mm × 5-cm; 1.8-µm packing [L1](#)

Column temperature: 50°

Flow rate: 2 mL/min

Injection volume: 20 µL

Run time: NLT 1.8 times the retention time of atorvastatin

System suitability

Sample: *Standard solution*

[NOTE—The relative retention times for amlodipine and atorvastatin are 0.12 and 1.0, respectively.]

Suitability requirements

Tailing factor: NMT 1.5 for amlodipine

Relative standard deviation: NMT 2.0% for amlodipine and atorvastatin

Analysis

Samples: *Standard solution and Sample solution*

Calculate the percentage of the labeled amount of amlodipine ($C_{20}H_{25}ClN_2O_5$) dissolved:

$$\text{Result} = (r_U/r_S) \times C_S \times V \times (M_{r1}/M_{r2}) \times (1/L) \times 100$$

r_U = peak response of amlodipine from the *Sample solution*

r_S = peak response of amlodipine from the *Standard solution*

C_S = concentration of [USP Amlodipine Besylate RS](#) in the *Standard solution* (mg/mL)

V = volume of *Medium*, 900 mL

M_{r1} = molecular weight of amlodipine, 408.88

M_{r2} = molecular weight of amlodipine besylate, 567.05

L = label claim of amlodipine (mg/Tablet)

Calculate the percentage of the labeled amount of atorvastatin ($C_{33}H_{35}FN_2O_5$) dissolved:

$$\text{Result} = (r_U/r_S) \times C_S \times V \times [M \times (M_{r1}/M_{r2})] \times (1/L) \times 100$$

r_U = peak response of atorvastatin from the *Sample solution*

r_S = peak response of atorvastatin from the *Standard solution*

C_S = concentration of [USP Atorvastatin Calcium RS](#) in the *Standard solution* (mg/mL)

V = volume of *Medium*, 900 mL

M = number of moles of atorvastatin per mole of atorvastatin calcium, 2

M_{r1} = molecular weight of atorvastatin, ▲558.65▲ (CN 1-May-2024)

M_{r2} = molecular weight of atorvastatin calcium, ▲1155.36▲ (CN 1-May-2024)

L = label claim of atorvastatin (mg/Tablet)

Tolerances: NLT 80% (Q) of the labeled amount of amlodipine ($C_{20}H_{25}ClN_2O_5$) and atorvastatin ($C_{33}H_{35}FN_2O_5$) are dissolved.

- **UNIFORMITY OF DOSAGE UNITS (905):** Meet the requirements

IMPURITIES

• ORGANIC IMPURITIES RELATED TO AMLODIPINE

Buffer 1: Add 7 mL of [triethylamine](#) in 1000 mL of [water](#) and adjust with [phosphoric acid](#) to a pH of 2.5. Add 1.8 g of [tetrabutylammonium hydrogen sulfate](#) and mix well.

Solution A: [Methanol](#) and *Buffer 1* (40:60)

Solution B: [Acetonitrile](#), [methanol](#), and *Buffer 1* (40:40:20)

Mobile phase: See [Table 3](#).

Table 3

Time (min)	Solution A (%)	Solution B (%)
0	90	10
2	90	10
7	75	25
16	70	30
18	55	45
24	25	75
30	10	90
31	0	100
35	0	100
36	90	10
40	90	10

Buffer 2: Add 7 mL of [triethylamine](#) in 1000 mL of [water](#). Adjust with [phosphoric acid](#) to a pH of 3.0.

Diluent 1: [Methanol](#) and [water](#) (50:50)

Diluent 2: [Methanol](#) and *Buffer 2* (50:50)

Standard stock solution: 0.7 mg/mL of [USP Amlodipine Besylate RS](#) in *Diluent 2*, prepared as follows. Transfer a suitable amount of [USP Amlodipine Besylate RS](#) to a suitable volumetric flask and dissolve in a quantity of [methanol](#), about 20% of the volume of the flask. Dilute with *Diluent 2* to volume.

Standard solution 1: 5 µg/mL of [USP Amlodipine Related Compound A RS](#) in *Diluent 1*

Standard solution 2: 3.5 µg/mL of [USP Amlodipine Besylate RS](#) from *Standard stock solution* in *Diluent 2*

Sample solution: Nominally 0.5 mg/mL of amlodipine in *Diluent 2*, prepared as follows. Finely powder NLT 25 Tablets and transfer a portion of the powder, equivalent to 50 mg of amlodipine to a 100-mL volumetric flask. Add about 40 mL of [methanol](#), shake to disperse, and sonicate for 15 min. Add about 40 mL of *Buffer 2* and sonicate for another 10 min. Dilute with *Diluent 2* to volume, centrifuge, and use the supernatant. Pass a portion of the solution through a suitable filter of 0.22-µm pore size. Prepare this solution fresh.

Chromatographic system

(See [Chromatography \(621\), System Suitability](#).)

Mode: LC

Detector: UV 270 nm for amlodipine related compound A; 360 nm for all other impurities

Column: 2.1-mm × 15-cm; 1.8-µm packing [L1](#)

Column temperature: 40°

Flow rate: 0.3 mL/min

Injection volume: 5 µL

System suitability

Sample: *Standard solution 2*

Suitability requirements

Tailing factor: NMT 2.0

Relative standard deviation: NMT 5.0%

Analysis**Samples:** *Standard solution 1, Standard solution 2, and Sample solution*

Calculate the percentage of amlodipine related compound A in the portion of Tablets taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times (M_{r1}/M_{r2}) \times 100$$

 r_U = peak response of amlodipine related compound A from the *Sample solution* r_S = peak response of amlodipine related compound A from *Standard solution 1* C_S = concentration of [USP Amlodipine Related Compound A RS](#) in *Standard solution 1* (mg/mL) C_U = nominal concentration of amlodipine in the *Sample solution* (mg/mL) M_{r1} = molecular weight of amlodipine related compound A free base, 406.86 M_{r2} = molecular weight of amlodipine related compound A fumarate, 522.94

Calculate the percentage of atorvastatin–amlodipine adduct or any unspecified degradation product in the portion of Tablets taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times (M_{r1}/M_{r2}) \times (1/F) \times 100$$

 r_U = peak response of each degradation product from the *Sample solution* r_S = peak response of amlodipine from *Standard solution 2* C_S = concentration of [USP Amlodipine Besylate RS](#) in *Standard solution 2* (mg/mL) C_U = nominal concentration of amlodipine in the *Sample solution* (mg/mL) M_{r1} = molecular weight of amlodipine, 408.88 M_{r2} = molecular weight of amlodipine besylate, 567.05 F = relative response factor (see [Table 4](#))**Acceptance criteria:** See [Table 4](#). Disregard peaks at the relative retention times of 2.18, 2.47 (atorvastatin), and 2.79 min.**Table 4**

Name	Relative Retention Time	Relative Response Factor	Acceptance Criteria, NMT (%)
Amlodipine related compound A	0.59	—	0.50
Amlodipine	1.00	—	—
Atorvastatin–amlodipine adduct ^a	3.49	0.47	0.50
Any unspecified degradation product	—	1.0	0.20
Total degradation products for amlodipine	—	—	1.0

^a 3-Ethyl 5-methyl 4-(2-chlorophenyl)-2-[(2-[(3R,5R)-7-[2-(4-fluorophenyl)-5-isopropyl-3-phenyl-4-(phenylcarbamoyl)-1H-pyrrol-1-yl]-3,5-dihydroxyheptanamido)ethoxy)methyl]-6-methyl-1,4-dihydropyridine-3,5-dicarboxylate.

Change to read:• **ORGANIC IMPURITIES RELATED TO ATORVASTATIN**

Buffer 1: Dissolve 6.8 g of [potassium dihydrogen phosphate](#) in 1000 mL of [water](#) and adjust with dilute [phosphoric acid](#) (1 in 10) to a pH of 3.4.

Buffer 2: Dissolve 6.8 g of [potassium dihydrogen phosphate](#) in 1000 mL of [water](#) and adjust with [triethylamine](#) to a pH of 7.0.

Solution A: [Tetrahydrofuran](#), [acetonitrile](#), and *Buffer 1* (5:25:70)

Solution B: [Tetrahydrofuran](#), [acetonitrile](#), and *Buffer 2* (5:70:25)

Mobile phase: See [Table 5](#).

Table 5

Time (min)	Solution A (%)	Solution B (%)
0	85	15
30	75	25
70	40	60
75	25	75
80	25	75
85	85	15
90	85	15

Diluent: [Acetonitrile](#) and [water](#) (50:50)

System suitability solution: Heat a suitable amount of [USP Atorvastatin Calcium RS](#) at 60° for 1 h for degradation; 0.55 mg/mL of degraded [USP Atorvastatin Calcium RS](#), 3 µg/mL each of [USP Atorvastatin Related Compound A RS](#), [USP Atorvastatin Related Compound B RS](#), [USP Atorvastatin Related Compound C RS](#), and [USP Atorvastatin Related Compound H RS](#) in *Diluent*. Sonication may be necessary for complete dissolution.

Standard solution: 2.7 µg/mL of [USP Atorvastatin Calcium RS](#) in *Diluent*

Sample solution: Nominally 0.5 mg/mL of atorvastatin in *Diluent*, prepared as follows. Transfer an amount equivalent to 50 mg of atorvastatin from a portion of NLT 20 finely powdered Tablets to a 100-mL volumetric flask. Add about 10 mL of [acetonitrile](#), shake to disperse, and sonicate for 5 min. Add about 70 mL of *Diluent* and sonicate for another 20 min. Dilute with *Diluent* to volume and centrifuge. Prepare this solution fresh.

Chromatographic system

(See [Chromatography \(621\)](#), [System Suitability](#).)

Mode: LC

Detector: UV 246 nm

Column: 4.6-mm × 25-cm; 4-µm packing [L11](#)

Column temperature: 45°

Flow rate: 1.2 mL/min

Injection volume: 20 µL

System suitability

Samples: *System suitability solution* and *Standard solution*

Suitability requirements

Resolution: NLT 1.0 between atorvastatin pyrrolidone analog and atorvastatin related compound A, *System suitability solution*

Relative standard deviation: NMT 5.0%, *Standard solution*

Analysis

Samples: *Standard solution* and *Sample solution*

Calculate the percentage of each atorvastatin specified or unspecified degradation product in the portion of Tablets taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times [M \times (M_{r1}/M_{r2})] \times (1/F) \times 100$$

r_U = peak response of each atorvastatin degradation product from the *Sample solution*

r_S = peak response of atorvastatin from the *Standard solution*

C_S = concentration of [USP Atorvastatin Calcium RS](#) in the *Standard solution* (mg/mL)

C_U = nominal concentration of atorvastatin in the *Sample solution* (mg/mL)

M = number of moles of atorvastatin per mole of atorvastatin calcium, 2

M_{r1} = molecular weight of atorvastatin, ▲558.65▲ (CN 1-May-2024)

M_{r2} = molecular weight of atorvastatin calcium, ▲1155.36▲ (CN 1-May-2024)

F = relative response factor (see [Table 6](#))

Acceptance criteria: See [Table 6](#). Disregard any impurity peaks less than 0.05% and the peaks from amlodipine related impurities.

Table 6

Name	Relative Retention Time	Relative Response Factor	Acceptance Criteria, NMT (%)
Atorvastatin pyrrolidone analog ^a	0.86	0.67	0.45
Atorvastatin related compound A ^b	0.91	—	—
Atorvastatin related compound B ^b	0.95	—	—
Atorvastatin	1.00	—	—
Atorvastatin related compound C ^b	1.04	—	—
Atorvastatin epoxy pyrrolooxazin 6-hydroxy analog ^c	1.35	0.39	0.5
Atorvastatin epoxy pyrrolooxazin 7-hydroxy analog ^d	1.40	0.52	0.5
Atorvastatin related compound H	1.78	1.0	1.0
Atorvastatin epoxy tetrahydrofuran analog ^e	1.96	0.63	0.5 ^g
Atorvastatin oxirane ^f	2.23	1.0	
Atorvastatin <i>tert</i> -butyl ester ^{b,h}	2.55	—	—
Any unspecified degradation product	—	—	0.20
Total degradation products for atorvastatin	—	—	2.0
Total degradation products ⁱ	—	—	3.0

^a (3*R*,5*R*)-7-[5-(4-Fluorophenyl)-3-isopropyl-2-oxo-4-phenyl-3-(phenylcarbamoyl)-2,3-dihydro-1*H*-pyrrol-1-yl]-3,5-dihydroxyheptanoic acid.

^b Process impurity included in the table for identification only. Process impurities are controlled in the drug substance, and are not to be reported or included in the total impurities for the drug product.

- c 4-{6-(4-Fluorophenyl)-7,8-epoxy-6-hydroxy-8a-isopropyl-7-phenyl-8-(phenylcarbamoyl)hexahydro-2H-pyrrolo[2,1-b][1,3]oxazin-2-yl}-3-hydroxybutanoic acid.
- d (3R)-4-(1b-(4-Fluorophenyl)-7-hydroxy-7-isopropyl-1a-phenyl-7a-(phenylcarbamoyl)hexahydro-1aH-oxireno[2',3':3,4]pyrrolo[2,1-b][1,3]oxazin-3-yl)-3-hydroxybutanoic acid.
- e 4-(4-Fluorophenyl)-2,4-dihydroxy-2-isopropyl-N,5-diphenyl-3,6-dioxabicyclo[3.1.0]hexane-1-carboxamide.
- f 3-(4-Fluorobenzoyl)-2-isobutyl-N,3-diphenyloxirane-2-carboxamide.
- g Sum of atorvastatin epoxy tetrahydrofuran analog and atorvastatin oxirane.
- h (3R,5R)-tert-Butyl 7-(2-(4-fluorophenyl)-5-isopropyl-3-phenyl-4-(phenylcarbamoyl)-1H-pyrrol-1-yl)-3,5-dihydroxyheptanoate.
- i Sum of the total degradation products for amlodipine from the test for *Organic Impurities Related to Amlodipine* and the total degradation products for atorvastatin from the test for *Organic Impurities Related to Atorvastatin*.

ADDITIONAL REQUIREMENTS

- **PACKAGING AND STORAGE:** Preserve in well-closed containers. Store at controlled room temperature.
- **LABELING:** When more than one *Dissolution* test is given, the labeling states the *Dissolution* test used only if *Test 1* is not used.

Change to read:

- **USP REFERENCE STANDARDS (11).**

[USP Amlodipine Besylate RS](#)

[USP Amlodipine Related Compound A RS](#)

3-Ethyl 5-methyl [2-(2-aminoethoxymethyl)-4-(2-chlorophenyl)-6-methyl-3,5-pyridinedicarboxylate] fumarate.

$C_{20}H_{23}ClN_2O_5 \cdot C_4H_4O_4$ 522.94

[USP Atorvastatin Calcium RS](#)

[USP Atorvastatin Related Compound A RS](#)

Calcium (3R,5R)-7-[2-isopropyl-4,5-diphenyl-3-(phenylcarbamoyl)-1H-pyrrol-1-yl]-3,5-dihydroxyheptanoate (1:2).

$C_{66}H_{70}CaN_4O_{10}$ 1119.38

[USP Atorvastatin Related Compound B RS](#)

Calcium (3S,5R)-7-[2-(4-fluorophenyl)-5-isopropyl-3-phenyl-4-(phenylcarbamoyl)-1H-pyrrol-1-yl]-3,5-dihydroxyheptanoate (1:2); also known as (3S,5R)-7-[3-(Phenylcarbamoyl)-5-(4-fluorophenyl)-2-isopropyl-4-phenyl-1H-pyrrol-1-yl]-3,5-dihydroxyheptanoic acid calcium salt.

$C_{66}H_{68}CaF_2N_4O_{10}$ ▲1155.36▲ (CN 1-May-2024)

[USP Atorvastatin Related Compound C RS](#)

Calcium (3R,5R)-7-[2,3-Bis(4-fluorophenyl)-5-isopropyl-4-(phenylcarbamoyl)-1H-pyrrol-1-yl]-3,5-dihydroxyheptanoate (1:2).

$C_{66}H_{66}CaF_4N_4O_{10}$ 1191.34

[USP Atorvastatin Related Compound H RS](#)

5-(4-Fluorophenyl)-1-{2-[(2R,4R)-4-hydroxy-6-oxotetrahydro-2H-pyran-2-yl]ethyl}-2-isopropyl-N,4-diphenyl-1H-pyrrole-3-carboxamide.

$C_{33}H_{33}FN_2O_4$ 540.64

Auxiliary Information - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
AMLODIPINE AND ATORVASTATIN TABLETS	Documentary Standards Support	SM22020 Small Molecules 2
REFERENCE STANDARD SUPPORT	RS Technical Services RSTECH@usp.org	SM22020 Small Molecules 2

Chromatographic Database Information: [Chromatographic Database](#)

Most Recently Appeared In:

Pharmacopeial Forum: Volume No. PF 43(2)

Current DocID: GUID-26183518-D649-435D-8C26-045F7CD2B9FB_10_en-US

DOI: https://doi.org/10.31003/USPNF_M1069_10_01

DOI ref: [uiyfk](#)