Status: Currently Official on 17-Feb-2025
Official Date: Official as of 01-Nov-2019
Document Type: USP Monographs
Docld: GUID-507405CA-704B-4B71-A30A-B1C14D1D23C8_2_en-US
DOI: https://doi.org/10.31003/USPNF_M6430_02_01
DOI Ref: i6ihr

© 2025 USPC Do not distribute

Atropine Sulfate Ophthalmic Solution

DEFINITION

Atropine Sulfate Ophthalmic Solution is a sterile, aqueous solution of Atropine Sulfate. It contains NLT 93.0% and NMT 107.0% of the labeled amount of atropine sulfate monohydrate $[(C_{17}H_{24}NO_3)_2 \cdot H_2SO_4 \cdot H_2O]$. It may contain suitable stabilizers and antimicrobial agents.

IDENTIFICATION

• A. The retention time of the major peak of the Sample solution corresponds to that of the Standard solution, as obtained in the Assay.

Delete the following:

B. IDENTIFICATION TESTS—GENERAL (191), Chemical Identification Tests, Sulfate

Sample solution: Evaporate to dryness a quantity of Ophthalmic Solution. Prepare a solution from the residue that contains the equivalent of 50 mg of atropine sulfate/mL.

Acceptance criteria: Meets the requirements

Add the following:

▲ B. The UV spectrum of the major peak of the Sample solution corresponds to that of the Standard solution, as obtained in the Assay. (IRA 1-

Nov-2019)

ASSAY

Change to read:

• PROCEDURE

▲Buffer A: 6.8 g/L of <u>sodium acetate</u> in <u>water</u>. To each liter add 3.5 mL of <u>triethylamine</u> and 6.6 mL of <u>glacial acetic acid</u>. Adjust with <u>glacial acetic acid</u>, if needed, to a pH of 4.5.

Buffer B: 6.8 g/L of sodium acetate in water. To each liter add 4 mL of glacial acetic acid. Adjust with glacial acetic acid, if needed, to a pH of

Mobile phase: Methanol and Buffer A (15:85) **Diluent:** Methanol and Buffer B (15:85)

Standard solution: 0.5 mg/mL of USP Atropine Sulfate RS in Diluent

Sample solution: Nominally 0.5 mg/mL of atropine sulfate monohydrate from a volume of Ophthalmic Solution prepared as follows. Rinse a 100-mL volumetric flask with *Diluent*. Transfer about 33 mL of *Diluent* to the flask and then add a 5.0-mL aliquot of Ophthalmic Solution using a "to contain" pipet (see *Volumetric Apparatus* (31)) to the flask. Shake vigorously. Dilute with *Diluent* to volume. Additional shaking may be needed to obtain a uniform solution.

Chromatographic system

(See Chromatography (621), System Suitability.)

Mode: LC

Detector: UV 225 nm. For Identification B, use a diode array detector in the range of 190-400 nm.

Column: 4.6-mm × 15-cm; 5-µm packing L10

Column temperature: 40° Flow rate: 1.2 mL/min Injection volume: 20 µL

Run time: NLT 3 times the retention time of atropine

System suitability

Sample: Standard solution

[Note—See <u>Table 1</u> for the relative retention times.]

Suitability requirements Tailing factor: NMT 2.0

Relative standard deviation: NMT 1.0%

Analysis

Samples: Standard solution and Sample solution

Calculate the percentage of the labeled amount of atropine sulfate monohydrate $[(C_{17}H_{23}NO_3)_2 \cdot H_2SO_4 \cdot H_2O]$ in the portion of Ophthalmic Solution taken:

r,, = peak response of atropine from the Sample solution

 $r_{\rm s}$ = peak response of atropine from the Standard solution

C_s = concentration of <u>USP Atropine Sulfate RS</u> in the Standard solution (mg/mL)

 C_{II} = nominal concentration of atropine sulfate in the Sample solution (mg/mL)

 M_{c1} = molecular weight of atropine sulfate monohydrate, 694.84

 M_{r_2} = molecular weight of anhydrous atropine sulfate, 676.82 (IRA 1-Nov-2019)

Acceptance criteria: 93.0%-107.0%

IMPURITIES

Change to read:

• ORGANIC IMPURITIES

Buffer A, Buffer B, Mobile phase, Diluent, Standard solution, Sample solution, and Chromatographic system: Proceed as directed in the

System suitability solution: 0.005 mg/mL of <u>atropic acid</u> and 0.5 mg/mL of <u>USP Atropine Sulfate RS</u> in *Diluent*

System suitability

Samples: Standard solution and System suitability solution [Note—See <u>Table 1</u> for the relative response factors.]

Suitability requirements

Resolution: NLT 1.5 between atropic acid and atropine, System suitability solution

Relative standard deviation: NMT 1.0%, Standard solution

Analysis

Samples: Standard solution and Sample solution

Calculate the percentage of each specified and any unspecified degradation product in the portion of Ophthalmic Solution taken:

Result =
$$(r_{1}/r_{S}) \times (C_{S}/C_{1}) \times (1/F) \times (M_{r1}/M_{r2}) \times 100$$

 r_{ij} = peak response of each specified and any unspecified degradation product from the Sample solution

r_s = peak response of atropine from the Standard solution

 C_S = concentration of <u>USP Atropine Sulfate RS</u> in the Standard solution (mg/mL)

C, = nominal concentration of atropine sulfate in the Sample solution (mg/mL)

F = relative response factor (see <u>Table 1</u>)

 M_{c1} = molecular weight of atropine sulfate monohydrate, 694.84

 M_{r2} = molecular weight of anhydrous atropine sulfate, 676.82

Acceptance criteria: See <u>Table 1</u>.

Table 1

Name	Relative Retention Time	Relative Response Factor	Acceptance Criteria, NMT (%)
Tropic acid ^a	0.69	2.0	7.0
Atropic acid ^b	0.87	12.2	1.0
Atropine	1.0	-	-
Apoatropine [©]	2.1	4.3	1.0
Any individual unspecified degradation product	_	1.0	1.0
Total impurities	_	-	7.0

https://trungtamthuoc.com/

- ^a 3-Hydroxy-2-phenylpropanoic acid.
- b 2-Phenylacrylic acid.
- $^{\rm c}~$ (1*R*,3*r*,5*S*)-8-Methyl-8-azabicyclo[3.2.1]octan-3-yl 2-phenylacrylate. $_{\blacktriangle~(IRA~1-Nov-2019)}$

SPECIFIC TESTS

- **PH (791)**: 3.5-6.0
- STERILITY TESTS (71): Meets the requirements

ADDITIONAL REQUIREMENTS

- PACKAGING AND STORAGE: Preserve in tight containers, and store at controlled room temperature.
- USP REFERENCE STANDARDS (11)

USP Atropine Sulfate RS

Auxiliary Information - Please check for your question in the FAQs before contacting USP.

Topic/Question	Contact	Expert Committee
ATROPINE SULFATE OPHTHALMIC SOLUTION	Documentary Standards Support	SM42020 Small Molecules 4
REFERENCE STANDARD SUPPORT	RS Technical Services RSTECH@usp.org	SM42020 Small Molecules 4

Chromatographic Database Information: Chromatographic Database

Most Recently Appeared In: Pharmacopeial Forum: Volume No. 45(2)

Current DocID: GUID-507405CA-704B-4B71-A30A-B1C14D1D23C8_2_en-US

DOI: https://doi.org/10.31003/USPNF_M6430_02_01

DOI ref: i6ihr