

Status: Currently Official on 13-Feb-2025  
Official Date: Official as of 01-Aug-2017  
Document Type: USP Monographs  
DocId: GUID-6651B071-F93A-411C-84B3-1FE2111C1AF8\_1\_en-US  
DOI: https://doi.org/10.31003/USPNF\_M8108\_01\_01  
DOI Ref: 787mu

© 2025 USPC  
Do not distribute

# Benzocaine, Butamben, and Tetracaine Hydrochloride Topical Solution

**DEFINITION**  
Benzocaine, Butamben, and Tetracaine Hydrochloride Topical Solution contains NLT 90.0% and NMT 110.0% of the labeled amount of benzocaine (C<sub>9</sub>H<sub>11</sub>NO<sub>2</sub>), butamben (C<sub>11</sub>H<sub>15</sub>NO<sub>2</sub>), and tetracaine hydrochloride (C<sub>15</sub>H<sub>24</sub>N<sub>2</sub>O<sub>2</sub> · HCl).

- IDENTIFICATION**
- A.** The retention times of the major peaks of the *Sample solution* correspond to those of the *Standard solution*, as obtained in the Assay.
  - B.** The UV spectrum of the major peaks of the *Sample solution* corresponds to that of the *Standard solution*, as obtained in the Assay.

**ASSAY**

• **PROCEDURE**

**Solution A:** 0.1% [formic acid](#) in [water](#)

**Solution B:** 0.1% [formic acid](#) in acetonitrile

**Mobile phase:** See [Table 1](#).

Table 1

| Time (min) | Solution A (%) | Solution B (%) |
|------------|----------------|----------------|
| 0          | 91             | 9              |
| 2.5        | 50             | 50             |
| 3.9        | 50             | 50             |
| 4          | 91             | 9              |
| 5          | 91             | 9              |

**Diluent:** [Acetonitrile](#) and [water](#) (10:90)

**Standard stock solution A:** 1750 µg/mL of [USP Benzocaine RS](#) prepared as follows. Transfer a suitable amount of [USP Benzocaine RS](#) to a suitable volumetric flask and dissolve in 80% of the total volume of *Diluent*. [NOTE—Sonication for about 5 min may be necessary.] Dilute with *Diluent* to volume.

**Standard stock solution B:** 125 µg/mL each of [USP Butamben RS](#) and [USP Tetracaine Hydrochloride RS](#) prepared as follows. Transfer a suitable amount of [USP Butamben RS](#) and [USP Tetracaine Hydrochloride RS](#) to a suitable volumetric flask and dissolve in 80% of the total volume of *Diluent*. [NOTE—Sonication for about 10 min may be necessary.] Dilute with *Diluent* to volume.

**Standard solution:** 175 µg/mL of [USP Benzocaine RS](#) from *Standard stock solution A* and 25 µg/mL each of [USP Butamben RS](#) and [USP Tetracaine Hydrochloride RS](#) from *Standard stock solution B* diluted in *Diluent*

**Sample solution:** Nominally 175 µg/mL of benzocaine and 25 µg/mL each of butamben and tetracaine hydrochloride, prepared as follows. Transfer a suitable portion of Topical Solution to a suitable volumetric flask and dissolve in 80% of the total volume of *Diluent*. [NOTE—Sonication for about 10 min may be necessary.] Dilute with *Diluent* to volume.

**Chromatographic system**  
(See [Chromatography \(621\)](#), [System Suitability](#).)

**Mode:** LC

**Detector:** UV 300 nm. For *Identification B*, use a diode array detector in the range of 200–400 nm.

**Column:** 2.1-mm × 5-cm; 1.7-µm packing [L1](#)

**Flow rate:** 0.6 mL/min

**Injection volume:** 1 µL

**System suitability**

**Sample:** *Standard solution*

[NOTE—The relative retention times for benzocaine, tetracaine, and butamben are about 0.71, 0.74, and 1.0, respectively.]

#### Suitability requirements

**Resolution:** NLT 2.0 between benzocaine and tetracaine

**Relative standard deviation:** NMT 2.0% for each of the three analyte peaks

#### Analysis

**Samples:** *Standard solution* and *Sample solution*

Calculate the percentage of the labeled amount of benzocaine ( $C_9H_{11}NO_2$ ), butamben ( $C_{11}H_{15}NO_2$ ), and tetracaine hydrochloride ( $C_{15}H_{24}N_2O_2 \cdot HCl$ ) in the portion of Topical Solution taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times 100$$

$r_U$  = peak response of the corresponding analyte from the *Sample solution*

$r_S$  = peak response of the corresponding analyte from the *Standard solution*

$C_S$  = concentration of the corresponding Reference Standard in the *Standard solution* (µg/mL)

$C_U$  = nominal concentration of the corresponding analyte in the *Sample solution* (µg/mL)

**Acceptance criteria:** 90.0%–110.0%

#### PERFORMANCE TESTS

- **MINIMUM FILL (755):** Meets the requirements

#### IMPURITIES

##### • ORGANIC IMPURITIES

**Mobile phase, Diluent, and Chromatographic system:** Proceed as directed in the Assay.

**System suitability solution:** 10 µg/mL each of [USP Benzocaine RS](#), [USP Tetracaine Hydrochloride RS](#), [USP Butamben RS](#), and [USP Ethyl 4-Nitrobenzoate RS](#) in *Diluent*

**Standard solution:** 3.4 µg/mL each of [USP Benzocaine RS](#) and [USP Ethyl 4-Nitrobenzoate RS](#) and 1 µg/mL of [USP Tetracaine Hydrochloride RS](#) in *Diluent*

**Sample solution:** Nominally 1.68 mg/mL of benzocaine, 0.24 mg/mL of butamben, and 0.24 mg/mL of tetracaine prepared as follows.

Accurately weigh a suitable quantity of Topical Solution into a suitable volumetric flask, dissolve with 20% of the total volume of acetonitrile, and dilute with water to volume. [NOTE—Sonication for about 1 min may be necessary.]

#### System suitability

**Samples:** *System suitability solution* and *Standard solution*

[NOTE—See [Table 2](#) for relative retention times.]

#### Suitability requirements

**Resolution:** NLT 2 between butamben and ethyl 4-nitrobenzoate; NLT 2 between benzocaine and tetracaine, *System suitability solution*

**Relative standard deviation:** NMT 5.0% for each of the analyte peaks, *Standard solution*

#### Analysis

**Samples:** *Standard solution* and *Sample solution*

Calculate the percentage of 4-aminobenzoic acid in the portion of Topical Solution taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times (1/F) \times 100$$

$r_U$  = peak response of 4-aminobenzoic acid from the *Sample solution*

$r_S$  = peak response of benzocaine from the *Standard solution*

$C_S$  = concentration of [USP Benzocaine RS](#) in the *Standard solution* (µg/mL)

$C_U$  = nominal concentration of benzocaine in the *Sample solution* (µg/mL)

$F$  = relative response factor (see [Table 2](#))

Calculate the percentage of ethyl 4-nitrobenzoate in the portion of Topical Solution taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times 100$$

$r_U$  = peak response of ethyl 4-nitrobenzoate from the *Sample solution*

$r_S$  = peak response of ethyl 4-nitrobenzoate from the *Standard solution*

$C_S$  = concentration of [USP Ethyl 4-Nitrobenzoate RS](#) in the *Standard solution* (µg/mL)

$C_U$  = nominal concentration of benzocaine in the *Sample solution* (µg/mL)

Calculate the percentage of tetracaine related compound B and any individual unspecified degradation product in the portion of Topical Solution taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times (1/F) \times 100$$

- $r_U$  = peak response of tetracaine related compound B or any individual unspecified degradation product from the *Sample solution*
- $r_S$  = peak response of tetracaine from the *Standard solution*
- $C_S$  = concentration of [USP Tetracaine Hydrochloride RS](#) in the *Standard solution* (µg/mL)
- $C_U$  = nominal concentration of tetracaine hydrochloride in the *Sample solution* (µg/mL)
- $F$  = relative response factor (see [Table 2](#))

**Acceptance criteria:** See [Table 2](#). Disregard any impurity peaks less than 0.05%.

Table 2

| Name                                           | Relative Retention Time | Relative Response Factor | Acceptance Criteria, NMT (%) |
|------------------------------------------------|-------------------------|--------------------------|------------------------------|
| 4-Aminobenzoic acid                            | 0.15                    | 1.3                      | 0.3                          |
| Benzocaine                                     | 0.70                    | —                        | —                            |
| Tetracaine                                     | 0.74                    | —                        | —                            |
| Tetracaine related compound B <sup>a</sup>     | 0.93                    | 0.6                      | 0.4                          |
| Butamben                                       | 1.0                     | —                        | —                            |
| Ethyl 4-nitrobenzoate                          | 1.04                    | —                        | 0.2                          |
| Any individual unspecified degradation product | —                       | 1.0                      | 0.4                          |
| Total degradation products                     | —                       | —                        | 2.0                          |

<sup>a</sup> 4-(Butylamino)benzoic acid.

ADDITIONAL REQUIREMENTS

- **PACKAGING AND STORAGE:** Preserve in tight containers, and avoid freezing.
- **USP REFERENCE STANDARDS** (11).

[USP Benzocaine RS](#)

[USP Butamben RS](#)

[USP Ethyl 4-Nitrobenzoate RS](#)

Ethyl *p*-nitrobenzoate.

C<sub>9</sub>H<sub>9</sub>NO<sub>4</sub> 195.17

[USP Tetracaine Hydrochloride RS](#)

**Auxiliary Information** - Please [check for your question in the FAQs](#) before contacting USP.

| Topic/Question                                                      | Contact                                       | Expert Committee          |
|---------------------------------------------------------------------|-----------------------------------------------|---------------------------|
| BENZOCAINE, BUTAMBEN, AND TETRACAINE HYDROCHLORIDE TOPICAL SOLUTION | <a href="#">Documentary Standards Support</a> | SM52020 Small Molecules 5 |

**Chromatographic Database Information:** [Chromatographic Database](#)

**Most Recently Appeared In:**

Pharmacopeial Forum: Volume No. PF 42(3)

**Current DocID: GUID-6651B071-F93A-411C-84B3-1FE2111C1AF8\_1\_en-US**

**DOI: [https://doi.org/10.31003/USPNF\\_M8108\\_01\\_01](https://doi.org/10.31003/USPNF_M8108_01_01)**

**DOI ref: [787mu](#)**

OFFICIAL