


Status: Currently Official on 13-Feb-2025  
Official Date: Official as of 01-Dec-2022  
Document Type: NF Monographs  
DocId: GUID-88582D45-6B5D-444E-928F-33FE0B9896FB\_4\_en-US  
DOI: [https://doi.org/10.31003/USPNF\\_M8570\\_04\\_01](https://doi.org/10.31003/USPNF_M8570_04_01)  
DOI Ref: ql2j4

© 2025 USPC  
Do not distribute

# Benzyl Alcohol

Portions of the monograph text that are national *USP* text, and are not part of the harmonized text, are marked with symbols (◆) to specify this fact.



C<sub>7</sub>H<sub>8</sub>O                      108.14  
Phenylmethanol    CAS RN®: 100-51-6.

### DEFINITION

Benzyl Alcohol contains NLT 98.0% and NMT the equivalent of 100.5% of phenylmethanol (C<sub>7</sub>H<sub>8</sub>O).

### IDENTIFICATION

*Change to read:*

- ◆▲ (NF 1-Dec-2022) A. [SPECTROSCOPIC IDENTIFICATION TESTS \(197\), Infrared Spectroscopy](#): 197F. On undried specimen. ▲▲ (NF 1-Dec-2022)

### ASSAY

*Change to read:*

• PROCEDURE

**Phenolphthalein solution:** Dissolve 0.1 g of phenolphthalein in 80 mL of ▲[alcohol](#)▲ (NF 1-Dec-2022) and dilute with water to 100.0 mL. To test for sensitivity, add 100 mL of carbon dioxide-free water to 0.1 mL of the *Phenolphthalein solution*. The solution is colorless. NMT 0.2 mL of 0.02 M sodium hydroxide is required to change the color to pink.

**Sample:** 0.900 g

▲**Titrant:** [1 N sodium hydroxide VS](#)▲ (NF 1-Dec-2022)

**Analysis:** To the *Sample* add 15 mL of a freshly prepared mixture of dried pyridine and acetic anhydride (7:1), and heat under a reflux condenser on a ▲▲ (NF 1-Dec-2022) water bath for 30 min. Cool, and add 25 mL of water. Using 0.25 mL of *Phenolphthalein solution* as the indicator, titrate with ▲*Titrant*▲ (NF 1-Dec-2022) Carry out a blank titration.  
Calculate the percentage content of phenylmethanol (C<sub>7</sub>H<sub>8</sub>O):

Result = 10.81 × (n<sub>1</sub> - n<sub>2</sub>) ▲×◆N▲ (NF 1-Dec-2022) /m

n<sub>1</sub>        = ▲volume of *Titrant*▲ (NF 1-Dec-2022) used for the blank (mL)

n<sub>2</sub>        = ▲volume of *Titrant*▲ (NF 1-Dec-2022) used for the sample (mL)

▲◆N      = actual normality of *Titrant* (mEq/mL)◆▲ (NF 1-Dec-2022)

m        = amount of sample taken (g)

**Acceptance criteria:** 98.0%–100.5%

### IMPURITIES

- [FATS AND FIXED OILS \(401\), Procedures, Peroxide Value](#): NMT 5

• RESIDUE ON EVAPORATION

**Analysis:** After ensuring that the Benzyl Alcohol complies with the test for *Fats and Fixed Oils <401>, Procedures, Peroxide Value*, evaporate 10.0 g to dryness in a tared quartz or porcelain crucible or platinum dish on a hot plate at a temperature not exceeding 200°. Ensure that the Benzyl Alcohol does not boil during evaporation. Dry the residue on the hot plate for 1 h, and allow to cool in a desiccator.

**Acceptance criteria:** The residue weighs NMT 5 mg, corresponding to NMT 0.05%.

*Change to read:*

• **ORGANIC IMPURITIES, BENZALDEHYDE, AND OTHER RELATED SUBSTANCES**

**Sample solution:** Use the Benzyl Alcohol sample under examination.

**Standard solution A:** Dissolve 0.100 g of ethylbenzene in 10.0 mL of the *Sample solution*. Dilute 2.0 mL of this solution with the *Sample solution* to 20.0 mL.

**Standard solution B:** Dissolve 2.000 g of dicyclohexyl in 10.0 mL of the *Sample solution*. Dilute 2.0 mL of this solution with the *Sample solution* to 20.0 mL.

**Reference solution A** (for use in nonparenteral applications): Dissolve 0.750 g of benzaldehyde and 0.500 g of cyclohexylmethanol in the *Sample solution*, and dilute with the *Sample solution* to 25.0 mL. Add 1.0 mL of this solution to a mixture of 2.0 mL of *Standard solution A* and 3.0 mL of *Standard solution B*, and dilute with the *Sample solution* to 20.0 mL.

**Reference solution B** (for use in parenteral applications): Dissolve 0.250 g of benzaldehyde and 0.500 g of cyclohexylmethanol in the *Sample solution*, and dilute with the *Sample solution* to 25.0 mL. Add 1.0 mL of this solution to a mixture of 2.0 mL of *Standard solution A* and 2.0 mL of *Standard solution B*, and dilute with the *Sample solution* to 20.0 mL.

**Chromatographic system**

(See [Chromatography \(621\)](#), [System Suitability](#).)

**Mode:** GC

**Detector:** Flame ionization

**Carrier:** Helium, chromatography grade

**Carrier linear velocity:** 25 cm/s, at 50°

**Detector temperature:** 310°

**Column:** 30-m × 0.32-mm; 0.5-µm film thickness, G16

**Temperatures**

**Injector:** 200°

**Column:** See [Table 1](#).

**Table 1**

Initial Temperature (°)	Temperature Ramp (°/min)	Final Temperature (°)	Hold Time at Final Temperature (min)
50	5	220	35

**System suitability**

**Sample:** For nonparenteral applications, use *Reference solution A*. For parenteral applications, use *Reference solution B*.

[NOTE—The retention time of benzyl alcohol is about 26 min. The relative retention times for ethylbenzene, dicyclohexyl, benzaldehyde, cyclohexylmethanol, and benzyl alcohol are about 0.28, 0.59, 0.68, 0.71, and 1.0, respectively.]

**Injection volume:** 0.1 µL without air plug

**Suitability requirements**

**Sensitivity:** Adjust the sensitivity of the detector so that the height of the peak due to ethylbenzene is NLT 30% of the full scale of the recorder.

**Resolution:** NLT 3.0 between the peaks corresponding to benzaldehyde and cyclohexylmethanol

**Analysis**

**Samples:** *Sample solution* and *Reference solution A* for nonparenteral applications and *Reference solution B* for parenteral applications

**Acceptance criteria** (nonparenteral applications): If any peaks are present in the chromatogram obtained with the *Sample solution* that have the same retention times as the peaks due to ethylbenzene and dicyclohexyl, subtract the areas of any such peaks from the peak areas at these retention times in the chromatograms of *Reference solution A* or *Reference solution B* (corrected peak areas of ethylbenzene and dicyclohexyl). Any such peaks in the *Sample solution* should be included in the assessments for the total of other peaks.

In the chromatogram obtained with the *Sample solution*, the area of any peak corresponding to benzaldehyde is NMT the difference between the area of the peak due to benzaldehyde in the chromatogram obtained with *Reference solution A* (0.15%) and the area of the peak due to benzaldehyde in the chromatogram obtained with the *Sample solution*.

In the chromatogram obtained with the *Sample solution*, the area of any peak corresponding to cyclohexylmethanol is NMT the difference between the area of the peak due to cyclohexylmethanol in the chromatogram obtained with *Reference solution A* (0.10%) and the area of the peak due to cyclohexylmethanol in the chromatogram obtained with the *Sample solution*.

In the chromatogram obtained with the *Sample solution*, the sum of the areas of any peak with a ▲ (NF 1-Dec-2022) retention time less than that of benzyl alcohol and apart from the peaks due to benzaldehyde and cyclohexylmethanol is NMT four times the area of ethylbenzene in *Reference solution A*, corrected if necessary as described above (0.04%).

In the chromatogram obtained with the *Sample solution*, the sum of the areas of any peak with a ▲ (NF 1-Dec-2022) retention time greater than that of benzyl alcohol is NMT the area of dicyclohexyl in *Reference solution A*, corrected if necessary as described above (0.3%).

Disregard any peak with an area less than 0.01 times that of the peak due to ethylbenzene in the chromatogram of *Reference solution A*, corrected if necessary as described above.

**Acceptance criteria** (parenteral applications): If any peaks are present in the chromatogram obtained with the *Sample solution* that have the same retention times as the peaks due to ethylbenzene and dicyclohexyl, subtract the areas of any such peaks from the peak areas at these retention times in the chromatograms of *Reference solution A* or *Reference solution B* (corrected peak areas of ethylbenzene and dicyclohexyl). Any such peaks in the *Sample solution* should be included in the assessments for the total of other peaks.

In the chromatogram obtained with the *Sample solution*, the area of any peak corresponding to benzaldehyde is NMT the difference between the area of the peak due to benzaldehyde in the chromatogram obtained with *Reference solution B* (0.05%) and the area of the peak due to benzaldehyde in the chromatogram obtained with the *Sample solution*.

In the chromatogram obtained with the *Sample solution*, the area of any peak corresponding to cyclohexylmethanol is NMT the difference between the area of the peak due to cyclohexylmethanol in the chromatogram obtained with *Reference solution B* (0.10%) and the area of the peak due to cyclohexylmethanol in the chromatogram obtained with the *Sample solution*.

In the chromatogram obtained with the *Sample solution*, the sum of the areas of any peak with a ▲ (NF 1-Dec-2022) retention time less than that of benzyl alcohol and apart from the peaks due to benzaldehyde and cyclohexylmethanol is NMT two times the area of ethylbenzene in *Reference solution B*, corrected if necessary as described above (0.02%).

In the chromatogram obtained with the *Sample solution*, the sum of the areas of any peak with a ▲ (NF 1-Dec-2022) retention time greater than that of benzyl alcohol is NMT the area of dicyclohexyl in *Reference solution B*, corrected if necessary as described above (0.2%).

Disregard any peak with an area less than 0.01 times that of the peak due to ethylbenzene in the chromatogram of *Reference solution B*, corrected if necessary as described above.

## SPECIFIC TESTS

### Change to read:

#### • ACIDITY

**Phenolphthalein solution:** Prepare as directed in the Assay.

**Analysis:** To 10 mL of Benzyl Alcohol add 10 mL of ▲ [alcohol](#) (NF 1-Dec-2022) and 1 mL of *Phenolphthalein solution*.

**Acceptance criteria:** NMT ▲ 1.0 (NF 1-Dec-2022) mL of 0.1 ▲ (NF 1-Dec-2022) sodium hydroxide is required to change the color of the indicator to pink.

#### • ♦ CLARITY OF SOLUTION

[NOTE—The *Sample solution* is to be compared to *Reference suspension 1* in diffused daylight 5 min after preparation of *Reference suspension 1*.]

**Hydrazine solution:** Transfer 1.0 g of hydrazine sulfate to a 100-mL volumetric flask, dissolve in and dilute with water to volume, and mix. Allow to stand 4–6 h before use.

**Methenamine solution:** Transfer 2.5 g of methenamine to a 100-mL glass-stoppered flask, add 25.0 mL of water, insert the glass stopper, and mix to dissolve.

**Primary opalescent suspension:** [NOTE—This suspension is stable for 2 months, provided it is stored in a glass container free from surface defects. The suspension must not adhere to the glass and must be well mixed before use.] Transfer 25.0 mL of *Hydrazine solution* to the *Methenamine solution* in the 100-mL glass-stoppered flask. Mix, and allow to stand for 24 h.

**Opalescence standard:** [NOTE—This suspension should not be used beyond 24 h after preparation.] Transfer 15.0 mL of the *Primary opalescent suspension* to a 1000-mL volumetric flask, dilute with water to volume, and mix.

**Reference suspension 1:** Transfer 5.0 mL of the *Opalescence standard* to a 100-mL volumetric flask, and dilute with water to volume.

**Reference suspension 2:** Transfer 10.0 mL of the *Opalescence standard* to a second 100-mL volumetric flask, and dilute with water to volume.

**Sample solution:** Dissolve 2.0 g of Benzyl Alcohol in 60 mL of water.

**Analysis:** Transfer a sufficient portion of the *Sample solution* to a test tube of colorless, transparent, neutral glass with a flat base and an internal diameter of 15–25 mm, to obtain a depth of 40 mm. Similarly transfer portions of *Reference suspension 1*, *Reference suspension 2*, and water to separate matching test tubes. Compare the *Sample solution*, *Reference suspension 1*, *Reference suspension 2*, and water in diffused daylight, viewing vertically against a black background (see [Visual Comparison \(630\)](#)). [NOTE—The diffusion of light must be such that *Reference suspension 1* can readily be distinguished from water, and that *Reference suspension 2* can readily be distinguished from *Reference suspension 1*.]

**Acceptance criteria:** The *Sample solution* shows the same clarity as that of water, or its opalescence is not more pronounced than that of *Reference suspension 1*.

#### • ♦ COLOR OF SOLUTION

**Sample solution:** Use the *Sample solution* prepared in the test for *Clarity of Solution*.

**Analysis:** Transfer a sufficient portion of the *Sample solution* to a test tube of colorless, transparent, neutral glass with a flat base and an internal diameter of 15–25 mm, to obtain a depth of 40 mm. Similarly transfer a portion of water to a separate matching test tube. Compare the color of the *Sample solution* with that of water in diffused daylight, viewing vertically against a white background (see [Visual Comparison \(630\)](#)).

**Acceptance criteria:** The *Sample solution* has the color of water.

#### • [REFRACTIVE INDEX \(831\)](#): 1.538–1.541 at 20°

## ADDITIONAL REQUIREMENTS

- ◆ **LABELING:** Where Benzyl Alcohol is intended for use in the manufacture of parenteral applications, it is so labeled. ◆
- ◆ **PACKAGING AND STORAGE:** Preserve in tight containers, protected from light. ◆
- ◆ **USP REFERENCE STANDARDS (11).**  
[USP Benzyl Alcohol RS](#)
- ◆

**Auxiliary Information** - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
BENZYL ALCOHOL	<a href="#">Documentary Standards Support</a>	SE2020 Simple Excipients

**Chromatographic Database Information:** [Chromatographic Database](#)

**Most Recently Appeared In:**

Pharmacopeial Forum: Volume No. PF 35(3)

**Current DocID:** GUID-88582D45-6B5D-444E-928F-33FE0B9896FB\_4\_en-US

**DOI:** [https://doi.org/10.31003/USPNF\\_M8570\\_04\\_01](https://doi.org/10.31003/USPNF_M8570_04_01)

**DOI ref:** [ql2j4](#)

OFFICIAL