

Status: Currently Official on 14-Feb-2025
Official Date: Official as of 01-May-2017
Document Type: USP Monographs
DocId: GUID-0B9EF5DF-DF00-4E90-8D92-42A2CA2F4BB7_1_en-US
DOI: https://doi.org/10.31003/USPNF_M9200_01_01
DOI Ref: n7kcx

© 2025 USPC
Do not distribute

Betamethasone Valerate Cream

DEFINITION

Betamethasone Valerate Cream contains an amount of betamethasone valerate ($C_{27}H_{37}FO_6$) equivalent to NLT 90.0% and NMT 110.0% of the labeled amount of betamethasone ($C_{22}H_{29}FO_5$), in a suitable cream base.

IDENTIFICATION

- A.** The retention time of the major peak of the *Sample solution* corresponds to that of the *Standard solution*, as obtained in the Assay.
- B.** The UV spectrum of the major peak of the *Sample solution* corresponds to that of the *Standard solution*, as obtained in the Assay.

ASSAY

• **PROCEDURE**

Solution A: Water

Solution B: [Acetonitrile](#)

Mobile phase: See [Table 1](#).

Table 1

Time (min)	Solution A (%)	Solution B (%)
0.0	63	37
7.0	63	37
15.0	30	70
19.0	30	70
19.1	10	90
21.0	10	90
21.1	63	37
25.0	63	37

Diluent A: [Tetrahydrofuran](#) and water (50:50)

Diluent B: [Acetonitrile](#) and water (40:60)

System suitability solution: 25 µg/mL of [USP Betamethasone Valerate RS](#) and 10 µg/mL of [USP Betamethasone Valerate Related Compound A RS](#) in *Diluent B*. Sonicate to dissolve if necessary.

Standard solution: 25 µg/mL of [USP Betamethasone Valerate RS](#) in *Diluent B*. Sonicate to dissolve if necessary.

Sample solution: Nominally 20 µg/mL of betamethasone, prepared as follows. Transfer 1.0 mg of betamethasone from a portion of Cream to a suitable glass centrifuge tube. Add 15.0 mL of *Diluent A* and mix with a vortex mixer to disperse the sample thoroughly. Add 35.0 mL of *Diluent B* and sonicate for 10 min with intermittent shaking. Centrifuge to obtain a clear supernatant. Pass through a suitable filter of 0.2-µm pore size using a glass syringe. Discard the first 1 mL.

Chromatographic system

(See [Chromatography \(621\)](#), [System Suitability](#).)

Mode: LC

Detector: UV 240 nm. For *Identification B*, use a diode array detector in the range of 200–400 nm.

Column: 4.6-mm × 15-cm; 3.5-µm packing [L1](#)

Temperatures

Autosampler: 4°

Column: Ambient

Flow rate: 1 mL/min

Injection volume: 100 µL

System suitability

Samples: *System suitability solution* and *Standard solution*

[NOTE—See [Table 2](#) for relative retention times.]

Suitability requirements

Resolution: NLT 2.0 between betamethasone valerate and betamethasone valerate related compound A, *System suitability solution*

Tailing factor: NMT 2.0, *Standard solution*

Relative standard deviation: NMT 1.0%, *Standard solution*

Analysis

Samples: *Standard solution* and *Sample solution*

Calculate the percentage of the labeled amount of betamethasone (C₂₂H₂₉FO₅) in the portion of Cream taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times (M_{r1}/M_{r2}) \times 100$$

r_U = peak response from the *Sample solution*

r_S = peak response from the *Standard solution*

C_S = concentration of [USP Betamethasone Valerate RS](#) in the *Standard solution* (µg/mL)

C_U = nominal concentration of betamethasone in the *Sample solution* (µg/mL)

M_{r1} = molecular weight of betamethasone, 392.46

M_{r2} = molecular weight of betamethasone valerate, 476.58

Acceptance criteria: 90.0%–110.0%

IMPURITIES

• ORGANIC IMPURITIES

Solution A, Solution B, Mobile phase, Diluent A, Diluent B, System suitability solution, Sample solution, and Chromatographic system: Proceed as directed in the Assay.

Standard solution: 0.25 µg/mL each of [USP Betamethasone RS](#), [USP Betamethasone Valerate RS](#), and [USP Betamethasone Valerate Related Compound A RS](#) in *Diluent B*. Sonicate to dissolve if necessary.

System suitability

Samples: *System suitability solution* and *Standard solution*

[NOTE—See [Table 2](#) for relative retention times.]

Suitability requirements

Resolution: NLT 2.0 between betamethasone valerate and betamethasone valerate related compound A, *System suitability solution*

Relative standard deviation: NMT 5%, *Standard solution*

Analysis

Samples: *Sample solution* and *Standard solution*

Calculate the percentage of each specified degradation product in the portion of Cream taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times 100$$

r_U = peak response of each specified degradation product from the *Sample solution*

r_S = peak response of the corresponding USP Reference Standard from the *Standard solution*

C_S = concentration of the corresponding USP Reference Standard in the *Standard solution* (µg/mL)

C_U = nominal concentration of betamethasone in the *Sample solution* (µg/mL)

Calculate the percentage of each unspecified degradation product in the portion of Cream taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times (M_{r1}/M_{r2}) \times 100$$

r_U = peak response of each unspecified degradation product from the *Sample solution*

r_S = peak response of betamethasone valerate from the *Standard solution*

C_S = concentration of [USP Betamethasone Valerate RS](#) in the *Standard solution* (µg/mL)

C_U = nominal concentration of betamethasone in the *Sample solution* (µg/mL)

M_{r1} = molecular weight of betamethasone, 392.46
 M_{r2} = molecular weight of betamethasone valerate, 476.58

Acceptance criteria: See [Table 2](#). Disregard any impurity peak less than 0.1%.

Table 2

Name	Relative Retention Time	Acceptance Criteria, NMT (%)
Betamethasone	0.30	1.0
Betamethasone valerate	1.00	—
Betamethasone valerate related compound A	1.04	1.0
Any individual unspecified degradation product	—	1.0
Total degradation products	—	2.0

SPECIFIC TESTS

- [MICROBIAL ENUMERATION TESTS \(61\)](#) and [TESTS FOR SPECIFIED MICROORGANISMS \(62\)](#): It meets the requirements of the tests for absence of *Staphylococcus aureus* and *Pseudomonas aeruginosa*.
- [MINIMUM FILL \(755\)](#): Meets the requirements

ADDITIONAL REQUIREMENTS

- **PACKAGING AND STORAGE:** Preserve in collapsible tubes or in tight containers.
- [USP REFERENCE STANDARDS \(11\)](#)
[USP Betamethasone RS](#)
[USP Betamethasone Valerate RS](#)
[USP Betamethasone Valerate Related Compound A RS](#)
9-Fluoro-11 β ,17-dihydroxy-16 β -methyl-3,20-dioxopregna-1,4-dien-21-yl valerate.
 $C_{27}H_{37}FO_6$ 476.58

Auxiliary Information - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
BETAMETHASONE VALERATE CREAM	Documentary Standards Support	SM52020 Small Molecules 5

Chromatographic Database Information: [Chromatographic Database](#)

Most Recently Appeared In:
Pharmacopeial Forum: Volume No. PF 41(5)

Current DocID: GUID-0B9EF5DF-DF00-4E90-8D92-42A2CA2F4BB7_1_en-US
DOI: https://doi.org/10.31003/USPNF_M9200_01_01
DOI ref: [n7kcx](#)