

Status: Currently Official on 14-Feb-2025
Official Date: Official Prior to 2013
Document Type: Reagents
DocId: GUID-2C9196F9-442E-4CD5-B5B9-CF9376341AE4_1_en-US
DOI: https://doi.org/10.31003/USPNF_R3115_01_01
DOI Ref: r1pug

© 2025 USPC
Do not distribute

Biuret Reagent TS

—Dissolve 1.5 g of cupric sulfate and 6.0 g of potassium sodium tartrate in 500 mL of water in a 1000-mL volumetric flask. Add 300 mL of carbonate-free sodium hydroxide solution (1 in 10), dilute with carbonate-free sodium hydroxide solution (1 in 10) to 1000 mL, and mix.

Auxiliary Information - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
BIURET REAGENT TS	Documentary Standards Support Associate Scientific Liaison.	HDQ Headquarters

Most Recently Appeared In:

Pharmacopeial Forum: Volume No. Information currently unavailable

Current DocID: [GUID-2C9196F9-442E-4CD5-B5B9-CF9376341AE4_1_en-US](#)

DOI: https://doi.org/10.31003/USPNF_R3115_01_01

DOI ref: [r1pug](#)

OFFICIAL