

Status: Currently Official on 14-Feb-2025  
Official Date: Official Prior to 2013  
Document Type: Reagents  
DocId: GUID-E841634D-8A0D-4FEC-B366-107DAC6AA749\_1\_en-US  
DOI: [https://doi.org/10.31003/USPNF\\_R1722\\_01\\_01](https://doi.org/10.31003/USPNF_R1722_01_01)  
DOI Ref: c9jcy

© 2025 USPC  
Do not distribute

## Blood

(for carbon monoxide test in gases)—Use *oxalated* or *defibrinated blood* of dogs, sheep, cattle, or human beings within 24 hours after bleeding. Prepare *oxalated blood* by adding 10 mg of sodium oxalate to each mL of the freshly drawn blood.

**Auxiliary Information** - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
BLOOD	<a href="#">Margareth R.C. Marques</a> Principal Scientific Liaison	HDQ Headquarters

**Most Recently Appeared In:**

Pharmacopeial Forum: Volume No. Information currently unavailable

**Current DocID:** [GUID-E841634D-8A0D-4FEC-B366-107DAC6AA749\\_1\\_en-US](#)

**DOI:** [https://doi.org/10.31003/USPNF\\_R1722\\_01\\_01](https://doi.org/10.31003/USPNF_R1722_01_01)

**DOI ref:** [c9jcy](#)

