


Status: Currently Official on 14-Feb-2025  
 Official Date: Official as of 01-May-2020  
 Document Type: NF Monographs  
 DocId: GUID-A1442242-9127-41FF-8C52-321A00ABBFF7\_2\_en-US  
 DOI: [https://doi.org/10.31003/USPNF\\_M11120\\_02\\_01](https://doi.org/10.31003/USPNF_M11120_02_01)  
 DOI Ref: vw91g

© 2025 USPC  
 Do not distribute

## Butylparaben

Portions of the monograph text that are national *USP* text, and are not part of the harmonized text, are marked with symbols (†) to specify this fact.



$C_{11}H_{14}O_3$  194.23  
 Benzoic acid, 4-hydroxy-, butyl ester;  
 Butyl *p*-hydroxybenzoate CAS RN®: 94-26-8.

### DEFINITION

Butylparaben contains NLT 98.0% and NMT 102.0% of  $C_{11}H_{14}O_3$ .

### IDENTIFICATION

**Change to read:**

- **A.** [▲SPECTROSCOPIC IDENTIFICATION TESTS \(197\), Infrared Spectroscopy: 197M▲](#) (CN 1-MAY-2020)
- **B.** [MELTING RANGE OR TEMPERATURE \(741\)](#): 68°–71°

### ASSAY

#### PROCEDURE

**Mobile phase, Sample solution, Standard solution B, and Chromatographic system:** Proceed as directed in the procedure for *Related Substances*.

#### System suitability

**Sample:** *Standard solution B*

#### Suitability requirements

**Relative standard deviation:** NMT 0.85% for six injections

#### Analysis

**Samples:** *Sample solution* and *Standard solution B*

Calculate the percentage of Butylparaben in the *Sample solution*:

$$\text{Result} = P \times (r_U \times C_S) / (r_S \times C_U)$$

$P$  = labeled purity of [USP Butylparaben RS](#) expressed as a percentage

$r_U$  = peak area of butylparaben from the *Sample solution*

$C_S$  = concentration of butylparaben in *Standard solution B* (mg/mL)

$r_S$  = peak area of butylparaben from *Standard solution B*

$C_U$  = concentration of Butylparaben in the *Sample solution* (mg/mL)

**Acceptance criteria:** 98.0%–102.0%

### IMPURITIES

- [RESIDUE ON IGNITION \(281\)](#): NMT 0.1%, determined on a 1.0-g sample

#### RELATED SUBSTANCES

**Mobile phase:** Methanol and a 6.8 g/L solution of potassium dihydrogen phosphate (1:1 v/v)

**Sample solution:** Dissolve 50.0 mg of Butylparaben in 2.5 mL of methanol, and dilute with *Mobile phase* to 50.0 mL. Dilute 10.0 mL of this solution with *Mobile phase* to 100.0 mL.

**Standard solution A:** 5.0 µg/mL each of *p*-hydroxybenzoic acid, [USP Propylparaben RS](#), and [USP Butylparaben RS](#) in *Mobile phase*

**Standard solution B:** Dissolve 50.0 mg of [USP Butylparaben RS](#) in 2.5 mL of methanol, and dilute with *Mobile phase* to 50.0 mL. Dilute 10.0 mL of this solution with *Mobile phase* to 100.0 mL.

**Standard solution C:** Dilute 1.0 mL of the *Sample solution* with *Mobile phase* to 20.0 mL. Dilute 1.0 mL of this solution with *Mobile phase* to 10.0 mL.

**Standard solution D:** 50 µg/mL of iso-butylparaben in *Mobile phase*

**Standard solution E:** *Standard solution D* in *Standard solution B* (1 in 100)

#### Chromatographic system

(See [Chromatography \(621\), System Suitability](#).)

**Mode:** LC

**Detector:** UV 272 nm

**Column:** 4.6-mm × 15-cm; 5-µm packing L1

**Column temperature:** 35°

**Flow rate:** 1.3 mL/min

**Injection volume:** 10 µL

**Run time:** About 1.5 times the retention time of butylparaben

#### System suitability

**Sample:** *Standard solutions A and E*

[NOTE—The retention time of butylparaben is about 22 min; the relative retention times for *p*-hydroxybenzoic acid, propylparaben, and iso-butylparaben with a reference to butylparaben are about 0.1, 0.5, and 0.9 min, respectively.]

#### Suitability requirements

**Resolution:** NLT 5.0 between the propylparaben and butylparaben peaks from *Standard solution A* and NLT 1.5 between the iso-butylparaben and butylparaben peaks from *Standard solution E*

#### Analysis

**Samples:** *Sample solution* and *Standard solution C*

[NOTE—Disregard any limit that is 0.2 times the area of the principal peak from *Standard solution C* (0.1%).]

#### Acceptance criteria

***p*-Hydroxybenzoic acid:** The peak area from the *Sample solution*, multiplied by 1.4 to correct for the calculation of content, is NMT the area of the principal peak from *Standard solution C* (0.5%).

**Unspecified impurities:** The peak area of each impurity from the *Sample solution* is NMT the area of the principal peak from *Standard solution C* (0.5%).

**Total impurities:** The total peak area for all impurities from the *Sample solution* is NMT twice the area of the principal peak from *Standard solution C* (1.0%).

#### SPECIFIC TESTS

• **Acidity:** To 2 mL of *Butylparaben solution* prepared in the *Color of Solution* test add 3 mL of alcohol, 5 mL of carbon dioxide-free water, and 0.1 mL of bromocresol green TS. Titrate with 0.10 N sodium hydroxide.

**Acceptance criteria:** NMT 0.1 mL is required to produce a blue color.

• **COLOR OF SOLUTION**

**Butylparaben solution:** Dissolve 1 g in alcohol, and dilute with alcohol to 10 mL.

**Acceptance criteria:** This solution is clear and not more intensely colored than alcohol or a solution prepared immediately before use by mixing 2.4 mL of ferric chloride CS, 1.0 mL of cobaltous chloride CS, and 0.4 mL of cupric sulfate CS with 0.3 N hydrochloric acid to make 10 mL, and diluting 5 mL of this solution with 0.3 N hydrochloric acid to make 100 mL. Make the comparison by viewing the solutions downward in matched color-comparison tubes against a white surface (see [Color and Achromicity \(631\)](#)).

#### ADDITIONAL REQUIREMENTS

• **PACKAGING AND STORAGE:** Preserve in well-closed containers.

• **USP REFERENCE STANDARDS (11).**

[USP Butylparaben RS](#)

[USP Propylparaben RS](#)

Auxiliary Information - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
BUTYLPARABEN	<a href="#">Documentary Standards Support</a>	SE2020 Simple Excipients

Chromatographic Database Information: [Chromatographic Database](#)

**Most Recently Appeared In:**

Pharmacopeial Forum: Volume No. PF 34(6)

**Current DocID: GUID-A1442242-9127-41FF-8C52-321A00ABBFF7\_2\_en-US**

**DOI: [https://doi.org/10.31003/USPNE\\_M11120\\_02\\_01](https://doi.org/10.31003/USPNE_M11120_02_01)**

**DOI ref: [vw91g](#)**

OFFICIAL