

Status: Currently Official on 16-Feb-2025
 Official Date: Official Prior to 2013
 Document Type: USP Monographs
 DocId: GUID-940B5132-0210-496D-BD35-4DC345CB36E5_1_en-US
 DOI: https://doi.org/10.31003/USPNF_M12255_01_01
 DOI Ref: hx8kk

© 2025 USPC
 Do not distribute

Camphorated Phenol Topical Solution

DEFINITION

Camphorated Phenol Topical Solution is a solution of Camphor and Phenol in eucalyptus oil and Light Mineral Oil. It contains NLT 90.0% and NMT 110.0% of the labeled amount of camphor ($C_{10}H_{16}O$) and phenol (C_6H_6O).

IDENTIFICATION

• **A.** The retention times of the camphor and phenol peaks of the *Sample solution* correspond to those of the *Standard solution*, as obtained in the Assay.

ASSAY

• PROCEDURE

Internal standard solution: 10 mg/mL of *n*-dodecane in chloroform

Standard stock solution: 9.6 mg/mL of [USP Phenol RS](#) and 22.4 mg/mL of [USP Camphor RS](#), in chloroform

Standard solution: 0.96 mg/mL of [USP Phenol RS](#), 2.24 mg/mL of [USP Camphor RS](#), and 1 mg/mL of *n*-dodecane in chloroform from the *Standard stock solution* and *Internal standard solution*

Sample solution: Transfer 1 g of Topical Solution to a 50-mL volumetric flask. Add 5.0 mL of *Internal standard solution*, and dilute with chloroform to volume.

Chromatographic system

(See [Chromatography \(621\)](#), [System Suitability](#).)

Mode: GC

Detector: Flame ionization

Column: 2-mm × 1.8-m glass; packed with 100- to 120-mesh S1A, coated with 15% G44

Carrier gas: Helium

Temperatures

Detector: 200°

Column: 140°

Injection volume: 1–2 µL

System suitability

Sample: *Standard solution*

[NOTE—The relative retention times are 0.3 for phenol, 0.8 for camphor, and 1.0 for the internal standard.]

Suitability requirements

Resolution: NLT 2.0 between camphor and the internal standard, and NLT 5.0 between phenol and camphor

Relative standard deviation: NMT 2.0% of the peak response ratio of camphor and phenol to the internal standard for five consecutive injections

Analysis

Samples: *Standard solution* and *Sample solution*

Calculate the percentage of the labeled amount of camphor (w/w) and phenol (w/w) in the portion of Topical Solution taken:

$$\text{Result} = (R_U/R_S) \times (C_S/C_U) \times 100$$

R_U = peak response ratio of the corresponding analyte to the internal standard from the *Sample solution*

R_S = peak response ratio of the corresponding analyte to the internal standard from the *Standard solution*

C_S = concentration of the appropriate USP Reference Standard in the *Standard solution* (mg/mL)

C_U = nominal concentration of the appropriate analyte in the *Sample solution* (mg/mL)

Acceptance criteria: 90.0%–110.0%

SPECIFIC TESTS

- [SPECIFIC GRAVITY \(841\)](#): 0.840–0.865

ADDITIONAL REQUIREMENTS

- **PACKAGING AND STORAGE:** Preserve in tight containers. Store at room temperature, avoid excessive heat, and close the cover after each use.
- [USP REFERENCE STANDARDS \(11\)](#).
[USP Camphor RS](#)
[USP Phenol RS](#)

Auxiliary Information - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
CAMPHORATED PHENOL TOPICAL SOLUTION	Documentary Standards Support	SM12020 Small Molecules 1
REFERENCE STANDARD SUPPORT	RS Technical Services RSTECH@usp.org	SM12020 Small Molecules 1

Chromatographic Database Information: [Chromatographic Database](#)

Most Recently Appeared In:

Pharmacopeial Forum: Volume No. PF 30(3)

Current DocID: [GUID-940B5132-0210-496D-BD35-4DC345CB36E5_1_en-US](#)

DOI: https://doi.org/10.31003/USPNF_M12255_01_01

DOI ref: [hx8kk](#)

OFFICIAL