

Status: Currently Official on 14-Feb-2025  
Official Date: Official Prior to 2013  
Document Type: USP Monographs  
DocId: GUID-D32532EA-9097-4AA5-BB33-8667DD0A074F\_1\_en-US  
DOI: [https://doi.org/10.31003/USPNF\\_M323\\_01\\_01](https://doi.org/10.31003/USPNF_M323_01_01)  
DOI Ref: v5v4x

© 2025 USPC  
Do not distribute

## Cephalexin Tablets for Oral Suspension

### DEFINITION

Cephalexin Tablets for Oral Suspension contain NLT 90.0% and NMT 110.0% of the labeled amount of cephalexin ( $C_{16}H_{17}N_3O_4S$ ).

### IDENTIFICATION

The retention time of the major peak of the *Sample solution* corresponds to that of the *Standard solution*, as obtained in the Assay.

### ASSAY

#### PROCEDURE

**Mobile phase:** 0.985 g/L of sodium 1-pentanesulfonate in a mixture of acetonitrile, methanol, triethylamine, and water (20:10:3:170), adjusted with phosphoric acid to a pH of  $3.0 \pm 0.1$

**Standard stock solution:** 1 mg/mL of [USP Cephalexin RS](#) in water

**Standard solution:** 0.4 mg/mL of cephalexin in *Mobile phase* from *Standard stock solution*

**Sample stock solution:** Nominally equivalent to 1 mg/mL of cephalexin from combined contents of NLT 20 powdered Tablets for Oral Suspension in water. Pass a portion of the solution through a filter having a 1- $\mu$ m or finer pore size.

**Sample solution:** 0.4 mg/mL of cephalexin in *Mobile phase* from *Sample stock solution*

#### Chromatographic system

(See [Chromatography \(621\)](#), [System Suitability](#).)

**Mode:** LC

**Detector:** UV 254 nm

**Column:** 4.6-mm  $\times$  25-cm; packing L1 of low acidity

**Flow rate:** 1.5 mL/min

**Injection size:** 20  $\mu$ L

#### System suitability

**Sample:** *Standard solution*

#### Suitability requirements

**Relative standard deviation:** NMT 2.0%

#### Analysis

**Samples:** *Standard solution* and *Sample solution*

Calculate the percentage of cephalexin ( $C_{16}H_{17}N_3O_4S$ ) in each Tablet for Oral Suspension:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times P \times F \times 100$$

$r_U$  = peak response from the *Sample solution*

$r_S$  = peak response from the *Standard solution*

$C_S$  = concentration of [USP Cephalexin RS](#) in the *Standard stock solution* (mg/mL)

$C_U$  = nominal concentration of cephalexin in the *Sample stock solution* (mg/mL)

$P$  = potency of cephalexin in [USP Cephalexin RS](#) ( $\mu$ g/mg)

$F$  = conversion factor, 0.001 mg/ $\mu$ g

**Acceptance criteria:** 90.0%–110.0%

### PERFORMANCE TESTS

• [DISINTEGRATION \(701\)](#): Tablets for Oral Suspension disintegrate in 3 min, using water at  $20 \pm 5^\circ$ .

• [DISSOLUTION \(711\)](#).

**Medium:** Water; 900 mL

**Apparatus 1:** Use 40-mesh cloth and 100 rpm.

**Time:** 30 min

**Standard solution:** 20  $\mu$ g/mL of [USP Cephalexin RS](#) in *Medium*

**Sample solution:** Pass a portion of the solution under test through a suitable filter. Dilute with *Medium*, if necessary, to a concentration of about 20 µg/mL.

**Spectrometric conditions**

(See [Ultraviolet-Visible Spectroscopy \(857\)](#).)

**Mode:** UV

**Analytical wavelength:** 262 nm

**Analysis**

**Samples:** *Standard solution* and *Sample solution*

**Tolerances:** NLT 80% (Q) of the labeled amount of cephalexin (C<sub>16</sub>H<sub>17</sub>N<sub>3</sub>O<sub>4</sub>S) is dissolved.

- **DISPERSION FINENESS:** Place 2 Tablets for Oral Suspension in 100 mL of water, and stir until completely dispersed. A smooth dispersion is obtained that passes through a No. 25 sieve.
- **UNIFORMITY OF DOSAGE UNITS (905):** Meets the requirements

**ADDITIONAL REQUIREMENTS**

- **PACKAGING AND STORAGE:** Preserve in tight containers at controlled room temperature.
- **USP REFERENCE STANDARDS (11):**  
[USP Cephalexin RS](#)

**Auxiliary Information** - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
CEPHALEXIN TABLETS FOR ORAL SUSPENSION	<a href="#">Documentary Standards Support</a>	SM12020 Small Molecules 1

**Chromatographic Database Information:** [Chromatographic Database](#)

**Most Recently Appeared In:**

Pharmacopeial Forum: Volume No. PF 36(1)

**Current DocID:** GUID-D32532EA-9097-4AA5-BB33-8667DD0A074F\_1\_en-US

**DOI:** [https://doi.org/10.31003/USPNF\\_M323\\_01\\_01](https://doi.org/10.31003/USPNF_M323_01_01)

**DOI ref:** [v5v4x](#)