

Status: Currently Official on 14-Feb-2025
 Official Date: Official Prior to 2013
 Document Type: USP Monographs
 DocId: GUID-3DA69C8D-C40E-4A1C-B502-03971136CC05_1_en-US
 DOI: https://doi.org/10.31003/USPNF_M16514_01_01
 DOI Ref: ghp35

© 2025 USPC
 Do not distribute

Chlorpheniramine Maleate and Pseudoephedrine Hydrochloride Oral Solution

» Chlorpheniramine Maleate and Pseudoephedrine Hydrochloride Oral Solution contains not less than 90.0 percent and not more than 110.0 percent of the labeled amounts of chlorpheniramine maleate ($C_{16}H_{19}ClN_2 \cdot C_4H_4O_4$) and pseudoephedrine hydrochloride ($C_{10}H_{15}NO \cdot HCl$).

Packaging and storage—Preserve in tight, light-resistant containers.

USP REFERENCE STANDARDS (11)—

[USP Chlorpheniramine Maleate RS](#)

[USP Pseudoephedrine Hydrochloride RS](#)

Identification—

A: The retention time of the major peak for chlorpheniramine maleate in the chromatogram of the *Assay preparation* corresponds to that of the *Standard preparation*, as obtained in the *Assay for chlorpheniramine maleate*.

B: The retention time of the major peak for pseudoephedrine hydrochloride in the chromatogram of the *Assay preparation* corresponds to that of the *Standard preparation* in the *Assay for pseudoephedrine hydrochloride*.

UNIFORMITY OF DOSAGE UNITS (905)—

FOR ORAL SOLUTION PACKAGED IN SINGLE-UNIT CONTAINERS: meets the requirements.

DELIVERABLE VOLUME (698)—

FOR ORAL SOLUTION PACKAGED IN MULTIPLE-UNIT CONTAINERS: meets the requirements.

Assay for chlorpheniramine maleate—

Mobile phase, System suitability solution, and Chromatographic system—Proceed as directed in the *Assay for chlorpheniramine maleate* under [Chlorpheniramine Maleate and Pseudoephedrine Hydrochloride Extended-Release Capsules](#).

Standard preparation—Dissolve an accurately weighed quantity of [USP Chlorpheniramine Maleate RS](#) in water to obtain a solution having a known concentration of about 1 mg per mL. Transfer 1.0 mL of this solution to a 100-mL volumetric flask, add 80 mL of *Mobile phase*, dilute with water to volume, and mix.

Assay preparation—Transfer an accurately measured volume of Oral Solution, equivalent to about 1 mg of chlorpheniramine maleate, to a 100-mL volumetric flask. Add about 80 mL of *Mobile phase*, dilute with water to volume, mix, and filter.

Procedure—Separately inject equal volumes (about 10 μ L) of the *Standard preparation* and the *Assay preparation* into the chromatograph, record the chromatograms, and measure the responses for the chlorpheniramine peaks. Calculate the quantity, in mg, of chlorpheniramine maleate ($C_{16}H_{19}ClN_2 \cdot C_4H_4O_4$) in the portion of Oral Solution taken by the formula:

$$(100C/V)(r_U/r_S)$$

in which *C* is the concentration, in mg per mL, of [USP Chlorpheniramine Maleate RS](#) in that *Standard preparation*; *V* is the volume, in mL, of Oral Solution taken for the *Assay preparation*; and r_U and r_S are the chlorpheniramine peak responses obtained from the *Assay preparation* and the *Standard preparation*, respectively.

Assay for pseudoephedrine hydrochloride—

Mobile phase, System suitability solution, and Chromatographic system—Proceed as directed in the *Assay for chlorpheniramine maleate* under [Chlorpheniramine Maleate and Pseudoephedrine Hydrochloride Extended-Release Capsules](#).

Standard preparation—Dissolve an accurately weighed quantity of [USP Pseudoephedrine Hydrochloride RS](#) in water to obtain a solution having a known concentration of about 1.5 mg per mL. Transfer about 1.0 mL of this solution to a 10-mL volumetric flask, add 8 mL of *Mobile phase*, dilute with water to volume, and mix.

Assay preparation—Transfer an accurately measured volume of Oral Solution, equivalent to about 15 mg of pseudoephedrine hydrochloride, to a 100-mL volumetric flask. Add 80 mL of *Mobile phase*, dilute with water to volume, mix, and filter.

Procedure—Separately inject equal volumes (about 10 μ L) of the *Standard preparation* and the *Assay preparation* into the chromatograph, record the chromatograms, and measure the responses for the pseudoephedrine peaks. Calculate the quantity, in mg, of pseudoephedrine

hydrochloride ($C_{10}H_{15}NO \cdot HCl$) in the portion of Oral Solution taken by the formula:

$$100(C/V)(r_U/r_S)$$

in which C is the concentration, in mg per mL, of [USP Pseudoephedrine Hydrochloride RS](#) in the *Standard preparation*; V is the volume, in mL, of Oral Solution taken for the *Assay preparation*; and r_U and r_S are the pseudoephedrine peak responses obtained from the *Assay preparation* and the *Standard preparation*, respectively.

Auxiliary Information - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
CHLORPHENIRAMINE MALEATE AND PSEUDOEPHEDRINE HYDROCHLORIDE ORAL SOLUTION	Documentary Standards Support	SM22020 Small Molecules 2

Chromatographic Database Information: [Chromatographic Database](#)

Most Recently Appeared In:

Pharmacopeial Forum: Volume No. PF 30(1)

Current DocID: [GUID-3DA69C8D-C40E-4A1C-B502-03971136CC05_1_en-US](#)

DOI: https://doi.org/10.31003/USPNF_M16514_01_01

DOI ref: [ghp35](#)

OFFICIAL