

Status: Currently Official on 14-Feb-2025
Official Date: Official Prior to 2013
Document Type: USP Monographs
DocId: GUID-E3703772-061F-4EDD-A60D-0E2BAC8419D5_1_en-US
DOI: https://doi.org/10.31003/USPNF_M18327_01_01
DOI Ref: 9pz0v

© 2025 USPC
Do not distribute

Clioquinol and Hydrocortisone Cream

DEFINITION

Clioquinol and Hydrocortisone Cream contains NLT 90.0% and NMT 110.0% of the labeled amounts of clioquinol (C_9H_5ClINO) and hydrocortisone ($C_{21}H_{30}O_5$) in a suitable cream base.

IDENTIFICATION

- **A.** The retention time of the clioquinol peak of the *Sample solution* corresponds to that of the *Standard solution*, as obtained in the Assay for *Clioquinol*.
- **B.** The retention time of the hydrocortisone peak of the *Sample solution* corresponds to that of the *Standard solution*, as obtained in the Assay for *Hydrocortisone*.

ASSAY

• CLIOQUINOL

Internal standard solution: 2 mg/mL of pyrene in pyridine

Standard stock solution: 3 mg/mL of [USP Clioquinol RS](#) in a mixture of pyridine and *n*-hexane (4:1)

Standard solution: Transfer 1.0 mL of the *Standard stock solution*, 1.0 mL of *N,O*-bis(trimethylsilyl)acetamide, and 1.0 mL of *Internal standard solution* to a suitable screw-capped glass vial fitted with a polytef-lined septum, and mix. Heat on a water bath at 50° for 15 min, and cool to room temperature.

Sample solution: Transfer nominally 150 mg of clioquinol from Cream to a 60-mL separator. Place the separator on its side in a vacuum oven at 45° for 4 h. Remove the separator, cool to room temperature, and add 15.0 mL of a mixture of pyridine and hexane (4:1). Insert the stopper in the separator, and mix until the specimen is completely dispersed. Transfer the contents of the separator to a 50-mL volumetric flask, rinse the separator with two 15-mL portions of a mixture of pyridine and hexane (4:1), collecting the rinsings in the volumetric flask, and dilute with the same solvent mixture to volume. Immediately transfer 1 mL of this solution to a dry, screw-capped glass vial, and evaporate with the aid of gentle heat and a stream of nitrogen to dryness. Dissolve the residue in 1.0 mL of a mixture of pyridine and hexane (4:1), add 1 mL each of *N,O*-bis(trimethylsilyl)acetamide and *Internal standard solution* to the screw-capped glass vial fitted with a polytef-lined septum, and mix. Heat on a water bath at 50° for 15 min, and cool to room temperature.

Chromatographic system

(See [Chromatography \(621\)](#), [System Suitability](#).)

Mode: GC

Detector: Flame ionization

Column: 2-mm × 1.8-m; packed with 3% liquid phase G3 on 80- to 100-mesh support SIAB

Temperatures

Column: 165°

Injection port: 170°

Detector: 250°

Carrier gas: Dry helium

Flow rate: 30 mL/min

Injection volume: 1 µL

System suitability

Sample: *Standard solution*

[NOTE—The relative retention times for clioquinol and pyrene are 0.6 and 1.0, respectively.]

Suitability requirements

Resolution: NLT 3.0 between the analyte and internal standard peaks

Relative standard deviation: NMT 2.0%

Analysis

Samples: *Standard solution* and *Sample solution*

Record the chromatograms to obtain NLT 40% of maximum recorder response, and measure the peak response of each component. Calculate the percentage of the labeled amount of clioquinol (C_9H_5ClINO) in the portion of Cream taken:

$$\text{Result} = (R_U/R_S) \times (C_S/C_U) \times 100$$

R_U = peak response ratio of clioquinol to the internal standard from the *Sample solution*

R_S = peak response ratio of clioquinol to the internal standard from the *Standard solution*

C_S = concentration of [USP Clioquinol RS](#) in the *Standard solution* (mg/mL)

C_U = nominal concentration of clioquinol in the *Sample solution* (mg/mL)

Acceptance criteria: 90.0%–110.0%

• HYDROCORTISONE

Mobile phase: Acetonitrile, methanol, and water (1:1:2.75)

Standard stock solution: 1 mg/mL of [USP Hydrocortisone RS](#) in alcohol

Standard solution: 100 µg/mL of [USP Hydrocortisone RS](#) in alcohol from the *Standard stock solution*

Sample solution: Transfer nominally 10 mg of hydrocortisone from Cream to a 50-mL centrifuge tube. Add 30 mL of alcohol, and heat on a steam bath just to boiling. Shake for 15 min, and centrifuge. Transfer the supernatant extract to a 100-mL volumetric flask. Repeat the extraction with two 20-mL portions of alcohol, combining the extracts in the 100-mL volumetric flask. Add alcohol to volume, mix, and filter.

System suitability stock solution: 0.5 mg/mL of methylparaben in alcohol

System suitability solution: Transfer 2 mL of *System suitability stock solution* and 20 mL of *Standard stock solution* into a 200-mL volumetric flask, and dilute with alcohol to volume.

Chromatographic system

(See [Chromatography \(621\)](#), [System Suitability](#).)

Mode: LC

Detector: UV 254 nm

Columns

Guard: Packing L2

Analytical: 3.9-mm × 30-cm; packing L1

Flow rate: 1 mL/min

Injection volume: 10 µL

System suitability

Sample: *System suitability solution*

[NOTE—The relative retention times for methylparaben and hydrocortisone are 0.6 and 1.0, respectively.]

Suitability requirements

Resolution: NLT 2.0 between the hydrocortisone and methylparaben peaks

Relative standard deviation: NMT 2.0%

Analysis

Samples: *Standard solution* and *Sample solution*

Calculate the percentage of the labeled amount of hydrocortisone ($C_{21}H_{30}O_5$) in the portion of Cream taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times 100$$

r_U = peak response from the *Sample solution*

r_S = peak response from the *Standard solution*

C_S = concentration of [USP Hydrocortisone RS](#) in the *Standard solution* (mg/mL)

C_U = nominal concentration of hydrocortisone in the *Sample solution* (mg/mL)

Acceptance criteria: 90.0%–110.0%

PERFORMANCE TESTS

- [MINIMUM FILL \(755\)](#): Meets the requirements

ADDITIONAL REQUIREMENTS

- **PACKAGING AND STORAGE:** Preserve in collapsible tubes or in tight, light-resistant containers.

- [USP REFERENCE STANDARDS \(11\)](#)
 - [USP Clioquinol RS](#)
 - [USP Hydrocortisone RS](#)

Auxiliary Information - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
CLIOQUINOL AND HYDROCORTISONE CREAM	Documentary Standards Support	SM12020 Small Molecules 1

Chromatographic Database Information: [Chromatographic Database](#)

Most Recently Appeared In:
Pharmacopeial Forum: Volume No. Information currently unavailable

Current DocID: GUID-E3703772-061F-4EDD-A60D-0E2BAC8419D5_1_en-US
DOI: https://doi.org/10.31003/USPNF_M18327_01_01
DOI ref: [9pz0v](#)

OFFICIAL