

Status: Currently Official on 14-Feb-2025  
Official Date: Official Prior to 2013  
Document Type: USP Monographs  
DocId: GUID-D8E0762D-4C3E-4597-8774-366F368CD47B\_1\_en-US  
DOI: [https://doi.org/10.31003/USPNF\\_M18935\\_01\\_01](https://doi.org/10.31003/USPNF_M18935_01_01)  
DOI Ref: 4ukw0

© 2025 USPC  
Do not distribute

## Clotrimazole and Betamethasone Dipropionate Cream

### DEFINITION

Clotrimazole and Betamethasone Dipropionate Cream contains NLT 90.0% and NMT 110.0% of the labeled amount of clotrimazole ( $C_{22}H_{17}ClN_2$ ) and an amount of betamethasone dipropionate equivalent to NLT 90.0% and NMT 110.0% of the labeled amount of betamethasone ( $C_{22}H_{29}FO_5$ ), in a suitable cream base.

### IDENTIFICATION

• **A.** The retention times of the major peaks for clotrimazole and betamethasone dipropionate of the *Sample solution* correspond to those of the *Standard solution*, as obtained in the Assay for clotrimazole and betamethasone.

### ASSAY

#### • PROCEDURE

**Buffer:** 6.6 g/L of dibasic ammonium phosphate in water

**Mobile phase:** Prepare a mixture of methanol and *Buffer* (7:3), and adjust with phosphoric acid to a pH of  $7.0 \pm 0.2$ . Pass through a membrane filter having a 0.45- $\mu$ m or finer pore size, and degas.

**Internal standard solution:** 0.15 mg/mL of progesterone in alcohol

**Clotrimazole stock solution:** 5 mg/mL of [USP Clotrimazole RS](#) in alcohol

**Betamethasone dipropionate stock solution:** 6.4J mg/mL of [USP Betamethasone Dipropionate RS](#) in alcohol, J being the ratio of the labeled amount of betamethasone (in mg/g) to the labeled amount of clotrimazole (in mg/g) in the Cream

**Clotrimazole related compound A stock solution:** 0.5 mg/mL of [USP Clotrimazole Related Compound A RS](#) in methanol

**Standard solution:** Transfer 1.0 mL of *Clotrimazole related compound A stock solution* to a suitable container, and evaporate to dryness in a water bath at room temperature under a stream of nitrogen. To the residue add 2.0 mL each of *Clotrimazole stock solution*, *Betamethasone dipropionate stock solution*, and *Internal standard solution*.

**Sample solution:** Weigh a portion of Cream equivalent to 10 mg of clotrimazole, and transfer to a screw-capped, 50-mL centrifuge tube. Add 2.0 mL of *Internal standard solution* and 4.0 mL of alcohol, place the cap on the tube, and heat at 60° in a water bath for 10 min, with occasional shaking. Remove the tube from the bath, cool in an ice bath for 20 min, and promptly centrifuge. Transfer a portion of the supernatant to a test tube, and use as the *Sample solution*.

#### Chromatographic system

(See [Chromatography \(621\)](#), [System Suitability](#).)

**Mode:** LC

**Detector:** UV 254 nm

**Column:** 4.6-mm  $\times$  25-cm; 10- $\mu$ m packing L1

**Flow rate:** 1.7 mL/min

**Injection volume:** 20  $\mu$ L

#### System suitability

**Sample:** *Standard solution*

[NOTE—The relative retention times for betamethasone dipropionate, clotrimazole related compound A, progesterone, and clotrimazole are about 1.0, 1.2, 1.4, and 1.7, respectively.]

#### Suitability requirements

**Resolution:** NLT 1.0 between betamethasone dipropionate and clotrimazole related compound A, NLT 1.5 between clotrimazole related compound A and progesterone, and NLT 1.8 between progesterone and clotrimazole

**Relative standard deviation:** NMT 2.0% determined from clotrimazole and betamethasone dipropionate and NMT 4.0% determined from clotrimazole related compound A

#### Analysis

**Samples:** *Standard solution* and *Sample solution*

Calculate the percentage of the labeled amount of clotrimazole ( $C_{22}H_{17}ClN_2$ ) in the portion of Cream taken:

$$\text{Result} = (R_U/R_S) \times (C_S/C_U) \times 100$$

$R_U$  = peak response ratio of clotrimazole to progesterone from the *Sample solution*

$R_S$  = peak response ratio of clotrimazole to progesterone from the *Standard solution*

$C_S$  = concentration of [USP Clotrimazole RS](#) in the *Standard solution* (mg/mL)

$C_U$  = nominal concentration of clotrimazole in the *Sample solution* (mg/mL)

Calculate the percentage of the labeled amount of betamethasone ( $C_{22}H_{29}FO_5$ ) in the portion of Cream taken:

$$\text{Result} = (R_U/R_S) \times (C_S/C_U) \times (M_{r1}/M_{r2}) \times 100$$

$R_U$  = peak response ratio of betamethasone dipropionate to progesterone from the *Sample solution*

$R_S$  = peak response ratio of betamethasone dipropionate to progesterone from the *Standard solution*

$C_S$  = concentration of [USP Betamethasone Dipropionate RS](#) in the *Standard solution* (mg/mL)

$C_U$  = nominal concentration of betamethasone in the *Sample solution* (mg/mL)

$M_{r1}$  = molecular weight of betamethasone, 392.46

$M_{r2}$  = molecular weight of betamethasone dipropionate, 504.60

**Acceptance criteria:** 90.0%–110.0% of the labeled amount of clotrimazole ( $C_{22}H_{17}ClN_2$ ); 90.0%–110.0% of the labeled amount of betamethasone ( $C_{22}H_{29}FO_5$ )

## IMPURITIES

### • ORGANIC IMPURITIES: LIMIT OF CLOTRIMAZOLE RELATED COMPOUND A

**Buffer, Mobile phase, Internal standard solution, Clotrimazole stock solution, Betamethasone dipropionate stock solution, Clotrimazole related compound A stock solution, Standard solution, Sample solution, Chromatographic system, and System suitability:** Use as directed in the Assay.

**Analysis:** Using the chromatograms of the *Standard solution* and *Sample solution* as obtained in the Assay, calculate the percentage of clotrimazole related compound A in the portion of Cream taken:

$$\text{Result} = (R_U/R_S) \times (C_S/C_U) \times 100$$

$R_U$  = peak response ratio of clotrimazole related compound A to progesterone from the *Sample solution*

$R_S$  = peak response ratio of clotrimazole related compound A to progesterone from the *Standard solution*

$C_S$  = concentration of [USP Clotrimazole Related Compound A RS](#) in the *Standard solution* (mg/mL)

$C_U$  = nominal concentration of clotrimazole in the *Sample solution* (mg/mL)

**Acceptance criteria:** NMT 5.0%

## PERFORMANCE TESTS

- [MINIMUM FILL \(755\)](#): Meets the requirements

## SPECIFIC TESTS

- [MICROBIAL ENUMERATION TESTS \(61\)](#) and [TESTS FOR SPECIFIED MICROORGANISMS \(62\)](#): It meets the requirements for the absence of *Staphylococcus aureus* and *Pseudomonas aeruginosa*.

## ADDITIONAL REQUIREMENTS

- **PACKAGING AND STORAGE:** Preserve in collapsible tubes or tight containers.

- [USP REFERENCE STANDARDS \(11\)](#).

[USP Betamethasone Dipropionate RS](#)

[USP Clotrimazole RS](#)

[USP Clotrimazole Related Compound A RS](#)

(*o*-Chlorophenyl)diphenylmethanol.

$C_{19}H_{15}ClO$

294.78

**Auxiliary Information** - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
CLOTRIMAZOLE AND BETAMETHASONE DIPROPIONATE CREAM	<a href="#">Documentary Standards Support</a>	SM12020 Small Molecules 1

**Chromatographic Database Information:** [Chromatographic Database](#)

**Most Recently Appeared In:**

Pharmacopeial Forum: Volume No. PF 29(3)

**Current DocID:** GUID-D8E0762D-4C3E-4597-8774-366F368CD47B\_1\_en-US

**DOI:** [https://doi.org/10.31003/USPNF\\_M18935\\_01\\_01](https://doi.org/10.31003/USPNF_M18935_01_01)

**DOI ref:** [4ukw0](#)

OFFICIAL