

Status: Currently Official on 14-Feb-2025
Official Date: Official Prior to 2013
Document Type: Reagents
DocId: GUID-7A9C71F1-5A82-4B1E-BBC0-7A2FC5AED6E3_1_en-US
DOI: https://doi.org/10.31003/USPNF_R3135_01_01
DOI Ref: iu2ii

© 2025 USPC
Do not distribute

Cobalt–Uranyl Acetate TS

–Dissolve, with warming, 40 g of uranyl acetate in a mixture of 30 g of glacial acetic acid and sufficient water to make 500 mL. Similarly, prepare a solution containing 200 g of cobaltous acetate in a mixture of 30 g of glacial acetic acid and sufficient water to make 500 mL. Mix the two solutions while still warm, and cool to 20°. Maintain the temperature at 20° for about 2 hours to separate the excess salts from solution, and then pass through a dry filter.

Auxiliary Information - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
COBALT–URANYL ACETATE TS	Margareth R.C. Marques Principal Scientific Liaison	HDQ Headquarters

Most Recently Appeared In:

Pharmacopeial Forum: Volume No. Information currently unavailable

Current DocID: [GUID-7A9C71F1-5A82-4B1E-BBC0-7A2FC5AED6E3_1_en-US](#)

DOI: https://doi.org/10.31003/USPNF_R3135_01_01

DOI ref: [iu2ii](#)

OFFICIAL