

Status: Currently Official on 14-Feb-2025  
Official Date: Official Prior to 2013  
Document Type: Reagents  
DocId: GUID-246151CA-F5BD-47FE-BBF8-62D15F0770B7\_1\_en-US  
DOI: [https://doi.org/10.31003/USPNF\\_R3144\\_01\\_01](https://doi.org/10.31003/USPNF_R3144_01_01)  
DOI Ref: c6ghz

© 2025 USPC  
Do not distribute

## Cupric Acetate TS, Stronger (*Barfoed's Reagent*)—

Dissolve 13.3 g of cupric acetate in a mixture of 195 mL of water and 5 mL of acetic acid.

**Auxiliary Information** - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
CUPRIC ACETATE TS, STRONGER (BARFOED'S REAGENT)	<a href="#">Margareth R.C. Marques</a> Principal Scientific Liaison	HDQ Headquarters

**Most Recently Appeared In:**

Pharmacopeial Forum: Volume No. Information currently unavailable

**Current DocID:** [GUID-246151CA-F5BD-47FE-BBF8-62D15F0770B7\\_1\\_en-US](#)

**DOI:** [https://doi.org/10.31003/USPNF\\_R3144\\_01\\_01](https://doi.org/10.31003/USPNF_R3144_01_01)

**DOI ref:** [c6ghz](#)

OFFICIAL