

Status: Currently Official on 14-Feb-2025
Official Date: Official Prior to 2013
Document Type: Reagents
DocId: GUID-DC0A98E1-49B6-4C46-AF20-2CC3506E18DF_1_en-US
DOI: https://doi.org/10.31003/USPNF_R3354_01_01
DOI Ref: b6ocq

© 2025 USPC
Do not distribute

Cupric Citrate TS 2, Alkaline

—With the aid of heat, dissolve about 173 g of sodium citrate dihydrate and 117 g of sodium carbonate monohydrate in about 700 mL of water, and filter. In a second flask, dissolve about 27.06 g of cupric sulfate ($\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$) in about 100 mL of water. Slowly combine the two solutions while stirring, and dilute with water to 1000 mL.

Auxiliary Information - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
CUPRIC CITRATE TS 2, ALKALINE	Margareth R.C. Marques Principal Scientific Liaison	HDQ Headquarters

Most Recently Appeared In:

Pharmacopeial Forum: Volume No. Information currently unavailable

Current DocID: [GUID-DC0A98E1-49B6-4C46-AF20-2CC3506E18DF_1_en-US](#)

DOI: https://doi.org/10.31003/USPNF_R3354_01_01

DOI ref: [b6ocq](#)

OFFICIAL