

Status: Currently Official on 14-Feb-2025
Official Date: Official Prior to 2013
Document Type: Reagents
DocId: GUID-CCED68BF-F36C-4212-994C-9A8B155A278B_1_en-US
DOI: https://doi.org/10.31003/USPNF_R3146_01_01
DOI Ref: 4ctu1

© 2025 USPC
Do not distribute

Cupric Citrate TS, Alkaline

—With the aid of heat, dissolve 173 g of dihydrated sodium citrate and 117 g of monohydrated sodium carbonate in about 700 mL of water, and filter through paper, if necessary, to obtain a clear solution. In a separate container dissolve 17.3 g of cupric sulfate in about 100 mL of water, and slowly add this solution, with constant stirring, to the first solution. Cool the mixture, add water to make 1000 mL, and mix.

Auxiliary Information - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
CUPRIC CITRATE TS, ALKALINE	Margareth R.C. Marques Principal Scientific Liaison	HDQ Headquarters

Most Recently Appeared In:

Pharmacopeial Forum: Volume No. Information currently unavailable

Current DocId: [GUID-CCED68BF-F36C-4212-994C-9A8B155A278B_1_en-US](#)

DOI: https://doi.org/10.31003/USPNF_R3146_01_01

DOI ref: [4ctu1](#)

OFFICIAL