

Status: Currently Official on 14-Feb-2025
 Official Date: Official as of 20-Jul-2022
 Document Type: USP Monographs
 DocId: GUID-212C5DF5-08FC-4F7F-9CDC-253952DDBDB8_3_en-US
 DOI: https://doi.org/10.31003/USPNF_M24530_03_01
 DOI Ref: b1nqg

© 2025 USPC
 Do not distribute

Diatrizoate Meglumine and Diatrizoate Sodium Solution

To view the Notice from the Expert Committee that posted in conjunction with this accelerated revision, please click
<https://www.uspnf.com/rb-diatrizoate-megulmine-diatrizoate-na-sol-20220729>.

Change to read:

DEFINITION

Diatrizoate Meglumine and Diatrizoate Sodium Solution is a solution of Diatrizoic Acid in Purified Water prepared with the aid of Meglumine and Sodium Hydroxide. ▲It can also be a solution of Diatrizoate Meglumine and Diatrizoate Sodium in Purified Water.▲ (RB 20-Jul-2022) It contains NLT 95.0% and NMT 105.0% of the labeled amounts of diatrizoate meglumine ($C_{11}H_{19}N_2O_4 \cdot C_7H_{17}NO_5$) and of iodine (I). It may contain small amounts of suitable buffers, Edetate Disodium, and flavoring agents.

IDENTIFICATION

• A. THIN-LAYER CHROMATOGRAPHIC IDENTIFICATION TEST (201)

Diluent: [Sodium hydroxide](#) in [methanol](#) (0.8 in 1000)

Standard solution: 1 mg/mL of [USP Diatrizoic Acid RS](#) in *Diluent*

Sample solution: 1 mg/mL from a volume of Solution, in *Diluent*

Developing solvent system: [Methanol](#), [chloroform](#), and [ammonium hydroxide](#) (10:20:2)

Analysis

Samples: *Sample solution* and *Standard solution*

Locate the spots using short-wavelength UV light.

Acceptance criteria: Meets the requirements

• B.

Sample solution: Nominally 500 mg of diatrizoate meglumine and diatrizoate sodium from a volume of Solution

Analysis: Evaporate the *Sample solution* to dryness and heat the residue so obtained in a crucible.

Acceptance criteria: Violet vapors are evolved.

ASSAY

• DIATRIZOATE MEGUMINE

Sample stock solution: 6 mg/mL of diatrizoate meglumine and diatrizoate sodium prepared as follows. Pipet a volume of Solution equivalent to 1.5 g of diatrizoate meglumine and diatrizoate sodium to a 250-mL volumetric flask, and dilute with [water](#) to volume.

Sample solution: Pipet 10 mL of the *Sample stock solution* into a glass-stoppered, 250-mL flask. Add 4 mL of 2 N [sulfuric acid](#) and 20 mL of [sodium metaperiodate](#) solution (1 in 200). Insert the stopper, and set aside in the dark for 1 h. Add 50 mL of [water](#), mix, and add 10 mL of [potassium iodide TS](#). Insert the stopper quickly and mix by swirling for 20 s.

Titrimetric system

Mode: Direct titration

Titrant: [0.1 N sodium thiosulfate VS](#)

Endpoint detection: Visual

Analysis: Immediately titrate the *Sample solution* with *Titrant*, using 3 mL of [starch TS](#). Perform a blank determination, and make any necessary correction. Each milliliter of 0.1 N sodium thiosulfate is equivalent to 10.11 mg of diatrizoate meglumine ($C_{11}H_{19}N_2O_4 \cdot C_7H_{17}NO_5$).

Acceptance criteria: 95.0%–105.0%

• IODINE

Sample stock solution: 0.08 g/mL of the total of diatrizoate meglumine and diatrizoate prepared as follows. Transfer an accurately measured volume of Solution, equivalent to 4 g of the total of diatrizoate meglumine and diatrizoate sodium, to a 50-mL volumetric flask, and dilute with [water](#) to volume, and mix.

Sample solution: Pipet 5 mL of *Sample stock solution* into a glass-stoppered, 125-mL conical flask. Add 30 mL of 1.25 N [sodium hydroxide](#) and 500 mg of powdered [zinc](#), connect the flask to a reflux condenser, and reflux the mixture for 1 h. Cool the flask to room temperature, rinse the condenser with 20 mL of [water](#), disconnect the flask from the condenser, and filter the mixture. Rinse the flask and filter thoroughly, adding the rinsings to the filtrate. Add 5 mL of [glacial acetic acid](#) and 1 mL of [tetrabromophenolphthalein ethyl ester TS](#).

Titrimetric system

Mode: Direct titration

Titrant: [0.05 N silver nitrate VS](#)

Endpoint detection: Visual

Analysis: Titrate the *Sample solution* with *Titrant* until the yellow precipitate just turns green. Each milliliter of 0.05 N silver nitrate is equivalent to 6.345 mg of iodine (I).

Acceptance criteria: 95.0%–105.0%

SPECIFIC TESTS

• **pH (791):** 6.0–7.6

• **IODINE AND IODIDE**

Solution A: [Sodium nitrite](#) solution (1 in 50)

Standard stock solution: [Potassium iodide](#) solution (1 in 4000)

Standard solution: Combine 2.0 mL of *Standard stock solution* and 22 mL of [water](#) in a stoppered 50-mL centrifuge tube.

Sample solution: Dilute a volume of *Solution*, equivalent to 2 g of diatrizoate meglumine and diatrizoate sodium, with [water](#) to 24 mL in a stoppered 50-mL centrifuge tube. Shake to promote dissolution.

Analysis

Samples: *Standard solution* and *Sample solution*

To each of the *Samples*, add 5 mL of [toluene](#) and 5 mL of 2 N [sulfuric acid](#), shake, and centrifuge. The toluene layer shows no red color.

Add 1 mL of *Solution A*, shake, and centrifuge.

Acceptance criteria: Any red color in the toluene layer of the *Sample solution* is not darker than the toluene layer of the *Standard solution*.

ADDITIONAL REQUIREMENTS

- **PACKAGING AND STORAGE:** Preserve in tight, light-resistant containers.
- **LABELING:** Label the container to indicate that the contents are not intended for parenteral use.
- **USP REFERENCE STANDARDS (11).**
[USP Diatrizoic Acid RS](#)

Auxiliary Information - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
DIATRIZOATE MEGLUMINE AND DIATRIZOATE SODIUM SOLUTION	Documentary Standards Support	SM42020 Small Molecules 4

Chromatographic Database Information: [Chromatographic Database](#)

Most Recently Appeared In:

Pharmacopeial Forum: Volume No. PF 27(6)

Current DocID: [GUID-212C5DF5-08FC-4F7F-9CDC-253952DDBDB8_3_en-US](#)

DOI: <https://doi.org/10.31003/USPNF.M24530.03.01>

DOI ref: [b1nqg](#)