

Status: Currently Official on 14-Feb-2025  
Official Date: Official Prior to 2013  
Document Type: Reagents  
DocId: GUID-DA52FE42-8766-40F5-A69F-C0A7C27693FA\_1\_en-US  
DOI: [https://doi.org/10.31003/USPNF\\_R3154\\_01\\_01](https://doi.org/10.31003/USPNF_R3154_01_01)  
DOI Ref: 9c6cj

© 2025 USPC  
Do not distribute

## Diazobenzenesulfonic Acid TS

—Place in a beaker 1.57 g of sulfanilic acid, previously dried at 105° for 3 hours, add 80 mL of water and 10 mL of diluted hydrochloric acid, and warm on a steam bath until dissolved. Cool to 15° (some of the sulfanilic acid may separate but will dissolve later), and add slowly, with constant stirring, 6.5 mL of sodium nitrite solution (1 in 10). Then dilute with water to 100 mL.

**Auxiliary Information** - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
DIAZOBENZENESULFONIC ACID TS	<a href="#">Margareth R.C. Marques</a> Principal Scientific Liaison	HDQ Headquarters

**Most Recently Appeared In:**

Pharmacopeial Forum: Volume No. Information currently unavailable

**Current DocId:** [GUID-DA52FE42-8766-40F5-A69F-C0A7C27693FA\\_1\\_en-US](#)

**DOI:** [https://doi.org/10.31003/USPNF\\_R3154\\_01\\_01](https://doi.org/10.31003/USPNF_R3154_01_01)

**DOI ref:** [9c6cj](#)

OFFICIAL