

Status: Currently Official on 14-Feb-2025
Official Date: Official Prior to 2013
Document Type: USP Monographs
DocId: GUID-AD9FDE73-0C5E-47EE-A8B6-096507A1E60A_2_en-US
DOI: https://doi.org/10.31003/USPNF_M25420_02_01
DOI Ref: gzk5x

© 2025 USPC
Do not distribute

Diethylcarbamazine Citrate Tablets

DEFINITION

Diethylcarbamazine Citrate Tablets contain NLT 95.0% and NMT 105.0% of the labeled amount of diethylcarbamazine citrate ($C_{10}H_{21}N_3O \cdot C_6H_8O_7$).

[NOTE—Diethylcarbamazine Citrate Tablets labeled solely for veterinary use are exempt from the requirements of the test for *Dissolution*.]

IDENTIFICATION

- **A. ~~IDENTIFICATION—ORGANIC NITROGENOUS BASES~~ (181):** Meet the requirements

ASSAY

PROCEDURE

Solution A: 10 g/L of [monobasic potassium phosphate](#) in [water](#)

Solution B: 31.24 mg/mL of [monobasic potassium phosphate](#) in [water](#)

Mobile phase: [Methanol](#) and *Solution A* (100:900)

Standard solution: 0.1 mg/mL of [USP Diethylcarbamazine Citrate RS](#) in *Solution B*

Sample solution: Nominally 0.1 mg/mL of diethylcarbamazine citrate prepared as follows. Transfer an amount of finely powdered Tablets (from NLT 20 Tablets) equivalent to 5 mg of diethylcarbamazine citrate to a 50-mL volumetric flask, and dissolve in and dilute with *Solution B* to volume.

Chromatographic system

Mode: LC

Detector: UV 220 nm

Column: 3.9-mm × 15-cm; 5-μm packing [L1](#)

Flow rate: 0.8 mL/min

Injection volume: 20 μL

System suitability

Sample: *Standard solution*

Suitability requirements

Relative standard deviation: NMT 2.0%

Analysis

Samples: *Standard solution* and *Sample solution*

Calculate the percentage of the labeled amount of diethylcarbamazine citrate ($C_{10}H_{21}N_3O \cdot C_6H_8O_7$) in the portion of Tablets taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times 100$$

r_U = peak response from the *Sample solution*

r_S = peak response from the *Standard solution*

C_S = concentration of [USP Diethylcarbamazine Citrate RS](#) in the *Standard solution* (mg/mL)

C_U = nominal concentration of diethylcarbamazine citrate in the *Sample solution* (mg/mL)

Acceptance criteria: 95.0%–105.0%

PERFORMANCE TESTS

- **~~DISINTEGRATION~~ (Z01):** (for Tablets labeled solely for veterinary use): 30 min

- **~~DISSOLUTION~~ (711):**

Procedure for a pooled sample

Medium: Water; 900 mL

Apparatus 2: 50 rpm

Time: 45 min

Buffer: 62.48 g/L of [monobasic potassium phosphate](#) in [water](#)

Sample solutions: Prepare by quantitatively diluting filtered portions of the solution under test with *Buffer* (1:1)

Analysis: Determine the amount of diethylcarbamazine citrate ($C_{10}H_{21}N_3O \cdot C_6H_8O_7$) dissolved as directed in the Assay.

Tolerances: NLT 75% (Q) of the labeled amount of diethylcarbamazine citrate ($C_{10}H_{21}N_3O \cdot C_6H_8O_7$) is dissolved.

- **UNIFORMITY OF DOSAGE UNITS (905):** Meet the requirements

IMPURITIES

• ORGANIC IMPURITIES

Solution A, Solution B, Mobile phase, and Chromatographic system: Proceed as directed in the Assay.

Citric acid solution: 2 mg/mL of citric acid in *Solution B*

Standard solution: 0.003 mg/mL of [USP Diethylcarbamazine Citrate RS](#) in *Solution B*

Sample solution: Transfer an amount of finely powdered Tablets (from NLT 20 Tablets) nominally equivalent to 300 mg of diethylcarbamazine citrate to a 100-mL volumetric flask. Dilute with *Solution B* to volume. Filter or centrifuge, and use the clear filtrate or supernatant.

Analysis

Samples: *Citric acid solution*, *Standard solution*, and *Sample solution*

Calculate the percentage of any individual impurity in the portion of Tablets taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times 100$$

r_U = peak response of any individual impurity from the *Sample solution*. [NOTE—Disregard any peak having a retention time corresponding to that of the main peak of the *Citric acid solution*.]

r_S = peak response of diethylcarbamazine citrate from the *Standard solution*

C_S = concentration of [USP Diethylcarbamazine Citrate RS](#) in the *Standard solution* (mg/mL)

C_U = nominal concentration of diethylcarbamazine citrate in the *Sample solution* (mg/mL)

Acceptance criteria: NMT 0.1%

ADDITIONAL REQUIREMENTS

- **PACKAGING AND STORAGE:** Preserve in tight containers.

- **USP REFERENCE STANDARDS (11).**

[USP Diethylcarbamazine Citrate RS](#)

Auxiliary Information - Please [check for your question in the FAQs](#) before contacting USP.

| Topic/Question | Contact | Expert Committee |
|------------------------------------|---|---------------------------|
| DIETHYLCARBAMAZINE CITRATE TABLETS | Documentary Standards Support | SM12020 Small Molecules 1 |

Chromatographic Database Information: [Chromatographic Database](#)

Most Recently Appeared In:

Pharmacopeial Forum: Volume No. PF 30(1)

Current DocID: GUID-AD9FDE73-0C5E-47EE-A8B6-096507A1E60A_2_en-US

DOI: https://doi.org/10.31003/USPNF_M25420_02_01

DOI ref: [gzk5x](#)