

Status: Currently Official on 17-Feb-2025

Official Date: Official as of 01-May-2020

Document Type: USP Monographs

DocId: GUID-296ADEC9-8E25-425F-8913-8E7EE37C3AC2_2_en-US


DOI: https://doi.org/10.31003/USPNF_M25660_02_01

DOI Ref: 8o3ez

© 2025 USPC

Do not distribute

Diethyltoluamide

 $C_{12}H_{17}NO$

191.27

Benzamide, *N,N*-diethyl-3-methyl-*N,N*-Diethyl-*m*-toluamide CAS RN®: 134-62-3; UNII: FB0C1XZV4Y.

DEFINITION

Diethyltoluamide contains NLT 95.0% and NMT 103.0% of the *meta*-isomer of diethyltoluamide ($C_{12}H_{17}NO$), calculated on the anhydrous basis.

IDENTIFICATION

Change to read:

- **A.** ▲ [SPECTROSCOPIC IDENTIFICATION TESTS \(197\)](#), [Infrared Spectroscopy: 197F](#) ▲ (CN 1-MAY-2020)
- **B.** The retention time of the major peak of the *Sample solution* corresponds to that of the *Standard solution*, as obtained in the Assay.

ASSAY

PROCEDURE

Solution A: 1 mL of phosphoric acid in 1000 mL of water**Mobile phase:** Methanol and *Solution A* (45:55)**Diluent:** Methanol and water (45:55)**System suitability solution:** 0.2 mg/mL of [USP Diethyltoluamide RS](#) (*m*-isomer) and 0.02 mg/mL of [USP Diethyltoluamide Related Compound A RS](#) in *Diluent*. [NOTE—[USP Diethyltoluamide Related Compound A RS](#) is the *p*-isomer of diethyltoluamide.]**Standard solution:** 0.2 mg/mL of [USP Diethyltoluamide RS](#) in *Diluent***Sample solution:** 0.2 mg/mL of Diethyltoluamide in *Diluent*

Chromatographic system

(See [Chromatography \(621\)](#), [System Suitability](#).)**Mode:** LC**Detector:** UV 235 nm**Column:** 4.6-mm × 15-cm; 3.5-μm packing L1**Flow rate:** 1 mL/min**Injection volume:** 10 μL

System suitability

Samples: *System suitability solution* and *Standard solution*

Suitability requirements

Resolution: NLT 1.5 between diethyltoluamide (*m*-isomer) and diethyltoluamide (*p*-isomer), *System suitability solution***Tailing factor:** NMT 2.0, *Standard solution***Relative standard deviation:** NMT 1.0%, *Standard solution*

Analysis

Samples: *Standard solution* and *Sample solution*Calculate the percentage of diethyltoluamide ($C_{12}H_{17}NO$) in the portion of Diethyltoluamide taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times 100$$

 r_U = peak response from the *Sample solution* r_S = peak response from the *Standard solution* C_S = concentration of the *Standard solution* (mg/mL) C_U = concentration of the *Sample solution* (mg/mL)

Acceptance criteria: 95.0%–103.0% on the anhydrous basis

SPECIFIC TESTS

- **SPECIFIC GRAVITY (841):** 0.996–1.002
- **REFRACTIVE INDEX (831):** 1.520–1.524
- **ACIDITY**

Sample solution: Dissolve 10.0 g in 50 mL of neutralized alcohol, and add phenolphthalein TS.

Analysis: Titrate the *Sample solution* with 0.010 N sodium hydroxide.

Acceptance criteria: NMT 4.0 mL of 0.010 N sodium hydroxide is required.

- **WATER DETERMINATION, Method I (921):** NMT 0.5%

ADDITIONAL REQUIREMENTS

- **PACKAGING AND STORAGE:** Preserve in tight containers.

- **USP REFERENCE STANDARDS (11).**

[USP Diethyltoluamide RS](#)

[USP Diethyltoluamide Related Compound A RS](#)

N,N-Diethyl-4-toluamide.

$C_{12}H_{17}NO$ 191.27

Auxiliary Information - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
DIETHYLTOLUAMIDE	Documentary Standards Support	SM12020 Small Molecules 1
REFERENCE STANDARD SUPPORT	RS Technical Services RSTECH@usp.org	SM12020 Small Molecules 1

Chromatographic Database Information: [Chromatographic Database](#)

Most Recently Appeared In:

Pharmacopeial Forum: Volume No. PF 39(2)

Current DocID: GUID-296ADEC9-8E25-425F-8913-8E7EE37C3AC2_2_en-US

DOI: https://doi.org/10.31003/USPNF_M25660_02_01

DOI ref: [8o3ez](#)