

Status: Currently Official on 14-Feb-2025
Official Date: Official as of 01-Dec-2016
Document Type: USP Monographs
DocId: GUID-B585D853-C2FF-4E25-9F35-1F72B50905E2_1_en-US
DOI: https://doi.org/10.31003/USPNF_M1603_01_01
DOI Ref: w2o17

© 2025 USPC
Do not distribute

Dolasetron Mesylate Compounded Oral Solution

DEFINITION

Dolasetron Mesylate Compounded Oral Solution contains NLT 90.0% and NMT 110.0% of the labeled amount of dolasetron mesylate ($C_{19}H_{20}N_2O_3 \cdot CH_4O_3S$).

Prepare Dolasetron Mesylate Compounded Oral Solution 10 mg/mL as follows (see [Pharmaceutical Compounding—Nonsterile Preparations \(795\)](#)).

Dolasetron Mesylate powder	1 g
Vehicle for Oral Solution (regular or sugar-free), <i>NF</i> , a sufficient quantity to make	100 mL

Add *Dolasetron Mesylate powder* and 15 mL of *Vehicle* to a mortar, and mix. Add the *Vehicle* in small portions almost to volume, and mix thoroughly after each addition. Transfer the contents of the mortar, stepwise and quantitatively, to a calibrated bottle. Add enough *Vehicle* to bring to final volume, and mix.

ASSAY

• **PROCEDURE**

Solution A: 0.05 M ammonium acetate adjusted with diluted ammonium hydroxide to a pH of 7.5

Mobile phase: Acetonitrile and *Solution A* (24:76). Filter, and degas.

Diluent: Acetonitrile and water (24:76)

Standard stock solution: 500 µg/mL of [USP Dolasetron Mesylate RS](#) in *Diluent*

Standard solution: 10 µg/mL of [USP Dolasetron Mesylate RS](#) from *Standard stock solution* in *Mobile phase*

Sample solution: 10 µg/mL of dolasetron mesylate prepared from Oral Solution and *Diluent*. Shake each sample thoroughly by hand for 15 s, centrifuge at 1000 rpm for 2 min, and use the supernatant.

Chromatographic system

(See [Chromatography \(621\), System Suitability](#).)

Mode: LC

Detector: UV 280 nm

Column: 4.6-mm × 15-cm; 3-µm packing L10

Column temperature: 30°

Flow rate: 0.8 mL/min

Injection volume: 5 µL

System suitability

Sample: *Standard solution*

[NOTE—The retention time for dolasetron mesylate is about 6.9 min.]

Suitability requirements

Relative standard deviation: NMT 1.4% for replicate injections

Analysis

Samples: *Standard solution* and *Sample solution*

Calculate the percentage of the labeled amount of dolasetron mesylate ($C_{19}H_{20}N_2O_3 \cdot CH_4O_3S$) in the portion of Oral Solution taken:

Result = $(r_U/r_S) \times (C_S/C_U) \times 100$

r_U = peak response from the *Sample solution*

r_S = peak response from the *Standard solution*

C_s = concentration of [USP Dolasetron Mesylate RS](#) in the *Standard solution* (µg/mL)

C_u = nominal concentration of dolasetron mesylate in the *Sample solution* (µg/mL)

Acceptance criteria: 90.0%–110.0%

SPECIFIC TESTS

- [pH \(791\)](#): 3.6–4.6

ADDITIONAL REQUIREMENTS

- **PACKAGING AND STORAGE:** Package in tight, light-resistant containers. Store at controlled room temperature, or in a refrigerator.
- **BEYOND-USE DATE:** NMT 90 days after the date on which it was compounded when stored at controlled room temperature, or in a refrigerator
- **LABELING:** Label it to state the *Beyond-Use Date*.
- **USP REFERENCE STANDARDS (11).**
[USP Dolasetron Mesylate RS](#)

Auxiliary Information - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
DOLASETRON MESYLATE COMPOUNDED ORAL SOLUTION	Brian Serumaga Science Program Manager	CMP2020 Compounding 2020

Chromatographic Database Information: [Chromatographic Database](#)

Most Recently Appeared In:
Pharmacopeial Forum: Volume No. PF 40(5)

Current DocID: [GUID-B585D853-C2FF-4E25-9F35-1F72B50905E2_1_en-US](#)
DOI: https://doi.org/10.31003/USPNF_M1603_01_01
DOI ref: [w2o17](#)

OFFICIAL