

Status: Currently Official on 14-Feb-2025
Official Date: Official as of 01-Sep-2020
Document Type: USP Monographs
DocId: GUID-062860E1-38C2-4F41-A4B1-ADE34B68DEDA_4_en-US
DOI: https://doi.org/10.31003/USPNF_M28020_04_01
DOI Ref: 9ne5f

© 2025 USPC
Do not distribute

Dopamine Hydrochloride Injection

To view the Notice from the Expert Committee that posted in conjunction with this accelerated revision, please click

<https://www.uspnf.com/rb/dopamine-hcl-inj-20200828>.

DEFINITION

Dopamine Hydrochloride Injection is a sterile solution of Dopamine Hydrochloride in Water for Injection. It contains NLT 95.0% and NMT 105.0% of the labeled amount of dopamine hydrochloride ($C_8H_{11}NO_2 \cdot HCl$). It may contain a suitable antioxidant.

[NOTE—Do not use the Injection if it is darker than slightly yellow or discolored in any other way.]

IDENTIFICATION

• **A. THIN-LAYER CHROMATOGRAPHIC IDENTIFICATION TEST (201).**

Standard solution: 1.6 mg/mL of [USP Dopamine Hydrochloride RS](#) in dilute [methanol](#) (1:5)

Sample solution: Nominally 1.6 mg/mL of dopamine hydrochloride prepared as follows. Transfer a volume of Injection to a suitable container, and dilute if necessary, with dilute [methanol](#) (1:5).

Chromatographic system

Application volume: 5 μ L

Developing solvent system: [n-Butyl alcohol](#), [glacial acetic acid](#), and [water](#) (4:1:1)

Analysis

Samples: *Standard solution* and *Sample solution*

Acceptance criteria: The R_f value of the principal spot from the *Sample solution* corresponds to that from the *Standard solution*.

ASSAY

• **PROCEDURE**

Solution A: 0.005 M [sodium 1-octanesulfonate](#) in 1% [glacial acetic acid](#)

Mobile phase: [Acetonitrile](#) and *Solution A*, (13:87). Filtered and degassed.

System suitability stock solution A: About 20 mg/mL of [benzoic acid](#) in [methanol](#)

System suitability stock solution B: About 5 mg/mL of [benzoic acid](#) from *System suitability stock solution A* prepared as follows. Dilute the *System suitability stock solution A* with *Mobile phase* (1:3, v/v).

Standard stock solution: About 1.6 mg/mL of [USP Dopamine Hydrochloride RS](#) in *Mobile phase*

System suitability solution: 0.16 mg/mL of [USP Dopamine Hydrochloride RS](#) and 0.5 mg/mL of [benzoic acid](#) prepared as follows. Transfer 10.0 mL of *System suitability stock solution B* and 10.0 mL of *Standard stock solution* to a 100-mL volumetric flask, dilute with *Mobile phase* to volume.

Standard solution: About 0.16 mg/mL of [USP Dopamine Hydrochloride RS](#) from the *Standard stock solution* in *Mobile phase*

Sample solution: Nominally 0.16 mg/mL of dopamine hydrochloride prepared as follows. Transfer an accurately measured volume of Injection, equivalent to about 16 mg of dopamine hydrochloride, to a 100-mL volumetric flask, dilute with *Mobile phase* to volume.

Chromatographic system

(See [Chromatography \(621\)](#), *System suitability*.)

Mode: LC

Detector: UV 280 nm

Column: 4-mm \times 30-cm; packing [L1](#)

Flow rate: 1.5 mL/min

Injection volume: 40 μ L

System suitability

Samples: *System suitability solution* and *Standard solution*

Suitability requirements

Resolution: NLT 4.0 between benzoic acid and dopamine hydrochloride, *System suitability solution*

Relative standard deviation: NMT 3.0%, *Standard solution*

Analysis

Samples: *Standard solution* and *Sample solution*

Calculate the percentage of the labeled amount of dopamine hydrochloride ($C_8H_{11}NO_2 \cdot HCl$) in the portion of Injection taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times 100$$

- r_U = peak response of dopamine from the *Sample solution*
- r_S = peak response of dopamine from the *Standard solution*
- C_S = concentration of [USP Dopamine Hydrochloride RS](#) in the *Standard solution* (mg/mL)
- C_U = nominal concentration of dopamine hydrochloride in the *Sample solution* (mg/mL)

Acceptance criteria: 95.0%–105.0%

SPECIFIC TESTS

- [BACTERIAL ENDOTOXINS TEST \(85\)](#): NMT 16.67 USP Endotoxin Units/mg of dopamine hydrochloride
- [PARTICULATE MATTER IN INJECTIONS \(788\)](#): Meets the requirements for small-volume injections
- [pH \(791\)](#): 2.5–5.0
- [OTHER REQUIREMENTS](#): It meets the requirements in [Injections and Implanted Drug Products \(1\)](#).

ADDITIONAL REQUIREMENTS

Change to read:

- **PACKAGING AND STORAGE:** Preserve in single-dose containers, ▲preferably▲ (RB 1-Sep-2020) of Type I glass.
- **LABELING:** Label it to indicate that the Injection is to be diluted with a suitable parenteral vehicle prior to intravenous infusion.
- [USP REFERENCE STANDARDS \(11\)](#)
[USP Dopamine Hydrochloride RS](#)

Auxiliary Information - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
DOPAMINE HYDROCHLORIDE INJECTION	Documentary Standards Support	SM22020 Small Molecules 2

Chromatographic Database Information: [Chromatographic Database](#)

Most Recently Appeared In:

Pharmacopeial Forum: Volume No. 47(5)

Current DocID: GUID-062860E1-38C2-4F41-A4B1-ADE34B68DEDA_4_en-US

DOI: <https://doi.org/10.31003/USPNF.M28020.04.01>

DOI ref: [9ne5f](#)