

Status: Currently Official on 14-Feb-2025
Official Date: Official as of 01-Jan-2020
Document Type: USP Monographs
DocId: GUID-939057CE-684F-421C-978A-E18FDA5522E3_2_en-US
DOI: https://doi.org/10.31003/USPNF_M28310_02_01
DOI Ref: j8fhv

© 2025 USPC
Do not distribute

Doxycycline Calcium Oral Suspension

DEFINITION

Doxycycline Calcium Oral Suspension is prepared from Doxycycline Hyclate and contains one or more suitable buffers, colors, diluents, flavors, and preservatives. It contains the equivalent of NLT 90.0% and NMT 125.0% of the labeled amount of doxycycline ($C_{22}H_{24}N_2O_8$).

IDENTIFICATION

- **A.** The UV spectrum of the major peak of the *Sample solution* corresponds to that of the *Standard solution*, as obtained in the Assay.
- **B.** The retention time of the major peak of the *Sample solution* corresponds to that of the *Standard solution*, as obtained in the Assay.

ASSAY

• **PROCEDURE**

Protect solutions containing doxycycline from light.

Solution A: Transfer 3.1 g of [monobasic potassium phosphate](#), 0.5 g of [edetate disodium](#), and 0.5 mL of [triethylamine](#) to a 1000-mL volumetric flask. Add about 850 mL of [water](#) and mix. Dilute with [water](#) to volume and adjust with [1 N sodium hydroxide](#) to a pH of 8.5 ± 0.1 . Pass through a suitable filter of 0.22- μ m pore size.

Solution B: [Methanol](#)

Mobile phase: See [Table 1](#).

Table 1

Time (min)	Solution A (%)	Solution B (%)
0.0	90	10
2.0	90	10
4.0	60	40
6.0	90	10
9.0	90	10

Diluent: [0.01 N hydrochloric acid](#)

Standard solution: 0.12 mg/mL of [USP Doxycycline Hyclate RS](#) in *Diluent*. Sonicate as needed to dissolve.

Sample solution: Nominally 0.1 mg/mL of doxycycline in *Diluent*, prepared as follows. Transfer an adequate amount of Oral Suspension, freshly mixed and free from air bubbles, to a suitable volumetric flask. Add 80% of the final volume of *Diluent*, sonicate for about 15 min, and dilute with *Diluent* to volume. Centrifuge a portion of the solution for 10 min at 3000 rpm and use the supernatant for analysis.

Chromatographic system

(See [Chromatography \(621\), System Suitability](#).)

Mode: LC

Detector: UV 270 nm. For *Identification A*, use a diode array detector in the range of 200-400 nm.

Column: 2.1-mm \times 5-cm; 1.7- μ m packing [L7](#)

[NOTE—A 1.7- μ m guard column with packing [L7](#) was used during method validation.]

Column temperature: 60°

Flow rate: 0.6 mL/min

Injection volume: 5 μ L

System suitability

Sample: *Standard solution*

Suitability requirements

Tailing factor: NMT 1.5

Relative standard deviation: NMT 2.0%

Analysis

Samples: *Standard solution* and *Sample solution*

Calculate the percentage of the labeled amount of doxycycline ($C_{22}H_{24}N_2O_8$) in the portion of Oral Suspension taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times P \times F \times 100$$

r_U = peak response from the *Sample solution*

r_S = peak response from the *Standard solution*

C_S = concentration of [USP Doxycycline Hyclate RS](#) in the *Standard solution* (mg/mL)

C_U = nominal concentration of doxycycline in the *Sample solution* (mg/mL)

P = potency of doxycycline in [USP Doxycycline Hyclate RS](#) (µg/mg)

F = conversion factor, 0.001 mg/µg

Acceptance criteria: 90.0%–125.0%

PERFORMANCE TESTS

- [UNIFORMITY OF DOSAGE UNITS \(905\)](#).

For single-unit containers

Acceptance criteria: Meets the requirements

- [DELIVERABLE VOLUME \(698\)](#): Meets the requirements

IMPURITIES

Delete the following:

▲ ORGANIC IMPURITIES

Protect solutions containing doxycycline from light.

Mobile phase, Diluent, and Chromatographic system: Proceed as directed in the Assay.

System suitability stock solution 1: 1 mg/mL each of [USP Doxycycline Related Compound A RS](#) and [USP Methacycline Hydrochloride RS](#) in *Diluent*

System suitability stock solution 2: 1.2 mg/mL of [USP Doxycycline Hyclate RS](#) in *Diluent*

System suitability solution: Transfer 5 mL of *System suitability stock solution 2* to a 25-mL volumetric flask, heat on a steam bath for 60 min, and evaporate to dryness on a hot plate, taking care not to char the residue. Dissolve the residue in *Diluent*, add 0.5 mL of *System suitability stock solution 1*, and dilute with *Diluent* to volume. Pass through a suitable filter of 0.20-µm pore size and use the filtrate. This solution contains a mixture of 4-epidoxycycline, doxycycline related compound A, methacycline, and doxycycline. [NOTE—The solution is stable up to 14 days when stored in a refrigerator.]

Standard solution: 2.3 µg/mL of [USP Doxycycline Hyclate RS](#) in *Diluent*

Sample solution: Nominally 2.0 mg/mL of doxycycline in *Diluent*, prepared as follows. Transfer an adequate amount of Oral Suspension, freshly mixed and free from air bubbles, to a suitable volumetric flask. Add 60% of the final volume of *Diluent*, sonicate for about 15 min, and dilute with *Diluent* to volume. Centrifuge a portion of the solution for 10 min at 3000 rpm and use the supernatant for analysis.

System suitability

Samples: *System suitability solution* and *Standard solution*

Suitability requirements

Resolution: NLT 1.5 between methacycline and 4-epidoxycycline; NLT 1.5 between 4-epidoxycycline and doxycycline related compound A; and NLT 2.0 between doxycycline related compound A and doxycycline, *System suitability solution*

Relative standard deviation: NMT 5.0% for doxycycline, *Standard solution*

Analysis

Samples: *Standard solution* and *Sample solution*

Calculate the percentage of each impurity in the portion of Oral Suspension taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times P \times F \times 100$$

r_U = peak response of each impurity from the *Sample solution*

r_S = peak response of doxycycline from the *Standard solution*

C_S = concentration of [USP Doxycycline Hyclate RS](#) in the *Standard solution* (mg/mL)

C_U = nominal concentration of doxycycline in the *Sample solution* (mg/mL)

P = potency of doxycycline in [USP Doxycycline Hyclate RS](#) (µg/mg)

F = conversion factor, 0.001 mg/µg

Acceptance criteria: See [Table 2](#). Disregard peaks less than 0.1%.

Table 2

Name	Relative Retention Time	Acceptance Criteria, NMT (%)
Methacycline ^{a,b}	0.64	—
4-Epidoxycycline ^c	0.79	0.5
Doxycycline related compound A (6-epidoxycycline) ^{b,d}	0.88	—
Doxycycline	1.0	—
Any individual unspecified impurity	—	0.5

^a (4S,4aR,5S,5aR,12aS)-4-(Dimethylamino)-1,4,4a,5,5a,6,11,12a-octahydro-3,5,10,12,12a-pentahydroxy-6-methylene-1,11-dioxo-2-naphthacenecarboxamide.

^b Process impurities that are controlled in the drug substance are not to be reported. They are listed here for information only.

^c (4R,4aR,5S,5aR,6R,12aS)-4-(Dimethylamino)-1,4,4a,5,5a,6,11,12a-octahydro-3,5,10,12,12a-pentahydroxy-6-methyl-1,11-dioxo-2-naphthacenecarboxamide.

^d (4S,4aR,5S,5aR,6S,12aS)-4-(Dimethylamino)-1,4,4a,5,5a,6,11,12a-octahydro-3,5,10,12,12a-pentahydroxy-6-methyl-1,11-dioxo-2-naphthacenecarboxamide.▲ (RB 1-Jan-2020)

SPECIFIC TESTS

- [pH \(791\)](#): 6.5–8.0

ADDITIONAL REQUIREMENTS

- **PACKAGING AND STORAGE:** Preserve in tight, light-resistant containers.

Change to read:

- [USP REFERENCE STANDARDS \(11\)](#)
[USP Doxycycline Hyclate RS](#)

▲ (RB 1-Jan-2020)

Auxiliary Information - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
DOXYCYCLINE CALCIUM ORAL SUSPENSION	Documentary Standards Support	SM12020 Small Molecules 1

Chromatographic Database Information: [Chromatographic Database](#)

Most Recently Appeared In:

Pharmacopeial Forum: Volume No. PF 42(4)

Current DocID: GUID-939057CE-684F-421C-978A-E18FDA5522E3_2_en-US

DOI: https://doi.org/10.31003/USPNF_M28310_02_01

DOI ref: [j8fhv](#)