


Status: Currently Official on 14-Feb-2025
Official Date: Official as of 01-May-2020
Document Type: USP Monographs
DocId: GUID-5B945100-1603-4D7D-A9C3-9B013A9ACD2F_5_en-US
DOI: https://doi.org/10.31003/USPNF_M29708_05_01
DOI Ref: m3ws

© 2025 USPC
Do not distribute

Eprinomectin



C₅₀H₇₅NO₁₄ (Component B_{1a}) 914.13
C₄₉H₇₃NO₁₄ (Component B_{1b}) 900.10
Component B_{1a}

Avermectin A_{1a}, 4''-(acetylamino)-5-O-demethyl-4''-deoxy-, (4''R)-;
(2aE,4E,5'S,6S,6'R,7S,8E,11R,13S,15S,17aR,20R,20aR,20bS)-6'--(S)-sec-butyl-5',6,6',7,10,11,14,15,17a,20,20a,20b-dodecahydro-20,20b-dihydroxy-5',6,8,19-tetramethyl-17-oxospiro[11,15-methano-2H,13H,17H-furo[4,3,2-pq][2,6]benzodioxacyclooctadecin-13,2'-[2H]pyran]-7-yl-4-O-(4-acetamido-2,4,6-trideoxy-3-O-methyl-α-L-lyxo-hexopyranosyl)-2,6-dideoxy-3-O-methyl-α-L-arabino-hexopyranoside [or (4''R)-4''-(acetylamino)-5-O-demethyl-4''-deoxyavermectin A_{1a}] CAS RN®: 133305-88-1; UNII: 00OY54D31C.

Component B_{1b}
Avermectin A_{1a}, 4''-(acetylamino)-5-O-demethyl-25-de(1-methylpropyl)-4''-deoxy-25-(1-methylethyl)-, (4''R)-;
(2aE,4E,5'S,6S,6'R,7S,8E,11R,13S,15S,17aR,20R,20aR,20bS)-5',6,6',7,10,11,14,15,17a,20,20a,20b-Dodecahydro-20,20b-dihydroxy-6'-isopropyl-5',6,8,19-tetramethyl-17-oxospiro[11,15-methano-2H,13H,17H-furo[4,3,2-pq][2,6]benzodioxacyclooctadecin-13,2'-[2H]pyran]-7-yl-4-O-(4-acetamido-2,4,6-trideoxy-3-O-methyl-α-L-lyxo-hexopyranosyl)-2,6-dideoxy-3-O-methyl-α-L-arabino-hexopyranoside [or (4''R)-4''-(acetylamino)-5-O-demethyl-25-de(1-methyl-propyl)-4''-deoxy-25-(1-methylethyl)avermectin A_{1a}] CAS RN®: 133305-89-2; UNII: 31OML2QZ0Q.

DEFINITION

Eprinomectin is a mixture of component B_{1a} (C₅₀H₇₅NO₁₄) and component B_{1b} (C₄₉H₇₃NO₁₄). It contains NLT 90.0% of component B_{1a} (C₅₀H₇₅NO₁₄) and NLT 95.0% of components B_{1a} (C₅₀H₇₅NO₁₄) and B_{1b} (C₄₉H₇₃NO₁₄), calculated on the anhydrous, solvent-free, and antioxidant-free basis. It may contain small amounts of a suitable antioxidant.

IDENTIFICATION

Change to read:

- **A.** **SPECTROSCOPIC IDENTIFICATION TESTS (197), Infrared Spectroscopy: 197M** (CN 1-MAY-2020)
- **B.** The retention times of the component B_{1a} peak and the component B_{1b} peak of the *Sample solution* correspond to those of the *Standard solution*, as obtained in the Assay.

ASSAY

• **PROCEDURE**

Solution A: 0.1% (v/v) solution of [perchloric acid](#) in [water](#)

Solution B: [Acetonitrile](#)

Mobile phase: See [Table 1](#).

Table 1

Time (min)	Solution A (%)	Solution B (%)
0	45	55

Time (min)	Solution A (%)	Solution B (%)
15	45	55
25	5	95
30	45	55
35	45	55

Diluent: [Methanol](#) and [water](#) (4:1)

Standard solution: 0.500 mg/mL of [USP Eprinomectin RS](#) in *Diluent*

System suitability solution: Transfer 4 mL of *Standard solution* to an LC vial. Add 2 drops of 1 M [sodium hydroxide](#) and let stand for 20 min prior to injection.

Sample solution: 0.500 mg/mL of Eprinomectin in *Diluent*

Chromatographic system

(See [Chromatography \(621\), System Suitability](#).)

Mode: LC

Detector: UV 245 nm

Column: 4.6-mm × 25-cm; 5-μm packing [L7](#)

Column temperature: 40°

Flow rate: 1.5 mL/min

Injection volume: 15 μL

System suitability

Samples: *Standard solution* and *System suitability solution*

[NOTE—For relative retention times, see [Table 2](#).]

Table 2

Components of the System Suitability Solution	Relative Retention Time
Impurity A	0.55
Component B _{1b}	0.77
Component B _{1a}	1.00
Impurities C + D	1.05
Impurity E	1.28

Suitability requirements

Resolution: NLT 3 between component B_{1b} and component B_{1a}; NLT 1 between component B_{1a} and impurities C + D, *System suitability solution*

Column efficiency: NLT 4,500 theoretical plates for component B_{1a}, *System suitability solution*

Symmetry factor: NMT 1.5 for component B_{1a}, *System suitability solution*

Relative standard deviation: NMT 1.0% from five injections for component B_{1a}, *Standard solution*

Analysis

Samples: *Standard solution* and *Sample solution*

Calculate the percentage of component B_{1a} (C₅₀H₇₅NO₁₄) in the portion of Eprinomectin taken:

$$\text{Result} = [r_{1a} / (r_{1a} + r_{1b})] \times 100$$

r_{1a} = peak area of component B_{1a} from the *Sample solution*

r_{1b} = peak area of component B_{1b} from the *Sample solution*

Calculate the percentage of component B_{1a} (C₅₀H₇₅NO₁₄) and component B_{1b} (C₄₉H₇₃NO₁₄) in the portion of Eprinomectin taken:

$$\text{Result} = (r_U / r_S) \times (C_S / C_U) \times 100$$

r_U = peak area of component B_{1a} or component B_{1b} from the *Sample solution*

r_S = peak area of component B_{1a} or component B_{1b} from the *Standard solution*

C_S = concentration of component B_{1a} or component B_{1b} in the *Standard solution* (mg/mL)

C_U = concentration of the *Sample solution* (mg/mL)

Acceptance criteria: NLT 90.0% of component B_{1a} and NLT 95.0% of components B_{1a} and B_{1b}, on the anhydrous, solvent-free, and antioxidant-free basis

IMPURITIES

• **RESIDUE ON IGNITION (281):** NMT 0.1%

• **LIMIT OF 8a-oxo-B_{1a}**

Solution A, Solution B, Diluent, Standard solution, System suitability solution, and Sample solution: Prepare as directed in the Assay.

Mobile phase: Acetonitrile and *Solution A* (13:7)

Butylated hydroxytoluene stock solution: 0.5 mg/mL of [butylated hydroxytoluene](#) in [methanol](#). Sonicate to dissolve, if necessary.

Butylated hydroxytoluene solution: 0.01 mg/mL of [butylated hydroxytoluene](#) from *Butylated hydroxytoluene stock solution* in *Diluent*

Chromatographic system

(See [Chromatography \(621\)](#), *System Suitability*.)

Mode: LC

Detector: UV 280 nm

Column: 4.6-mm × 25-cm; 5-μm packing [L7](#)

Column temperature: 40°

Flow rate: 1.5 mL/min

Injection volume: 15 μL

System suitability 1

Samples: *Standard solution* and *System suitability solution*

System suitability determination: Use the conditions as directed for *Chromatographic system* and the suitability requirements for *System suitability* as directed in the Assay.

System suitability 2

Samples: *Sample solution* and *Butylated hydroxytoluene solution*

Suitability requirements

Relative standard deviation: NMT 3.0% from six injections, *Butylated hydroxytoluene solution*

Analysis

Samples: *Sample solution* and *Butylated hydroxytoluene solution*

[NOTE—The retention time for 8a-oxo-B_{1a} is 4–9 min from the *Sample solution*, and the retention time for butylated hydroxytoluene is 12–17 min from *Butylated hydroxytoluene solution*.]

Calculate the percentage of 8a-oxo-B_{1a}, on the anhydrous, solvent-free, and antioxidant-free basis in the portion of Eprinomectin taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times F \times P \times 100$$

r_U = peak area of 8a-oxo-B_{1a} from the *Sample solution*

r_S = peak area of butylated hydroxytoluene from the *Butylated hydroxytoluene solution*

C_S = concentration of butylated hydroxytoluene in the *Butylated hydroxytoluene solution* (mg/mL)

C_U = concentration of Eprinomectin in the *Sample solution* (mg/mL)

F = relative response factor for butylated hydroxytoluene with respect to 8a-oxo-B_{1a}, 0.4

P = purity of butylated hydroxytoluene used to prepare the *Butylated hydroxytoluene solution*

Acceptance criteria: NMT 0.5%

Change to read:

• **ORGANIC IMPURITIES**

Solution A, Solution B, Mobile phase, Diluent, Standard solution, System suitability solution, Sample solution, Chromatographic system, and System suitability: Proceed as directed in the Assay.

Analysis

Sample: *Sample solution*

Calculate the percentage of each impurity in the portion of Eprinomectin taken:

$$\text{Result} = (r_U/r_T) \times 100$$

r_U = peak area of each individual related substance from the *Sample solution*

r_T = sum of the responses of all the peaks

- Acceptance criteria:** [NOTE—See [Table 2](#) for the relative retention times of impurity A and impurity E.]
- Impurities with relative retention times of 0.23, 0.93, and 1.16 with respect to the B_{1a} peak:** NMT 1.0%
- Impurity A:** NMT 1.0%
- Impurity E:** NMT 1.0%
- All other known impurities:** NMT 0.5%
- Total unknown impurities:** NMT 1.0%▲▲ (ERR 1-Jun-2019)
- Total impurities:** NMT 5.0%

SPECIFIC TESTS

- [OPTICAL ROTATION \(781S\)](#), [Procedures](#), [Specific Rotation](#)
Sample solution: 5 mg/mL of Eprinomectin in [chloroform](#)
Acceptance criteria: +132° to +140°, determined at 405 nm on the anhydrous, solvent-free, and antioxidant-free basis
- [WATER DETERMINATION \(921\)](#), [Method I](#), [Method Ia](#)
Sample: 0.250 g
Acceptance criteria: NMT 2.0%

ADDITIONAL REQUIREMENTS

- **PACKAGING AND STORAGE:** Preserve in tight containers, and store between 2° and 8° at ambient humidity.
- **LABELING:** Label it to state the name(s) and amount(s) of any added substance(s). Label to indicate that it is for veterinary use only.
- [USP REFERENCE STANDARDS \(11\)](#).
[USP Eprinomectin RS](#)

Auxiliary Information - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
EPRINOMECTIN	Documentary Standards Support	SM32020 Small Molecules 3

Chromatographic Database Information: [Chromatographic Database](#)

Most Recently Appeared In:
Pharmacopeial Forum: Volume No. PF 42(4)

Current DocID: GUID-5B945100-1603-4D7D-A9C3-9B013A9ACD2F_5_en-US
DOI: <https://doi.org/10.31003/USPNF.M29708.05.01>
DOI ref: [rn3ws](#)