

Status: Currently Official on 14-Feb-2025  
Official Date: Official Prior to 2013  
Document Type: Reagents  
DocId: GUID-3ADB0BFE-0311-4B1A-A20B-3EE0DF4747A0\_1\_en-US  
DOI: [https://doi.org/10.31003/USPNF\\_R1978\\_01\\_01](https://doi.org/10.31003/USPNF_R1978_01_01)  
DOI Ref: f jy1k

© 2025 USPC  
Do not distribute

## Equilenin,

$C_{18}H_{18}O_2$  266.33 CAS RN<sup>®</sup>: 517-09-9.—Colorless or white crystals or crystalline powder. Insoluble in water; soluble in chloroform and in dioxane; moderately soluble in alcohol.

- **MELTING RANGE, Class II (741):** between 256° and 260°.
- **SPECIFIC ROTATION (781):** between +85° and +88°, determined in a solution in dioxane containing 75 mg of equilenin in each 10 mL.
- **ABSORPTION MAXIMA:** An alcohol solution exhibits absorption maxima at 231, 282, 325, and 340 nm.

**Auxiliary Information** - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
EQUILENIN	<a href="#">Margareth R.C. Marques</a> Principal Scientific Liaison	HDQ Headquarters

**Most Recently Appeared In:**

Pharmacopeial Forum: Volume No. Information currently unavailable

**Current DocID:** GUID-3ADB0BFE-0311-4B1A-A20B-3EE0DF4747A0\_1\_en-US

**DOI:** [https://doi.org/10.31003/USPNF\\_R1978\\_01\\_01](https://doi.org/10.31003/USPNF_R1978_01_01)

**DOI ref:** [f jy1k](#)

OFFICIAL