

Status: Currently Official on 14-Feb-2025
Official Date: Official as of 01-May-2020
Document Type: USP Monographs
DocId: GUID-F63BC4D0-6225-4D78-9830-F891DEA4772E_4_en-US
DOI: https://doi.org/10.31003/USPNF_M10965_04_01
DOI Ref: ypk9e

© 2025 USPC
Do not distribute

Fentanyl Citrate Compounded Injection

DEFINITION

Fentanyl Citrate Compounded Injection contains NLT 90.0% and NMT 110.0% of the labeled amount of fentanyl ($C_{22}H_{28}N_2O$). It contains no bacteriostat or other preservative.

Prepare Fentanyl Citrate Compounded Injection, 50 mcg/mL, as follows (see [Pharmaceutical Compounding—Sterile Preparations \(797\)](#)).

Fentanyl (as fentanyl citrate)	5000 mcg (7870 mcg)
Sodium hydroxide solution or Hydrochloric acid solution	To adjust pH
Sodium Chloride for Injection (0.9%), a sufficient quantity to make	100 mL

Dissolve Fentanyl Citrate in sufficient Sodium Chloride for Injection (0.9%) to bring to final volume. Adjust with Sodium hydroxide solution or Hydrochloric acid solution to a pH of 4–7.5. Pass the solution through a suitable sterile filter of 0.22- μ m pore size into appropriate sterile syringes or container(s). [NOTE—Do not use a cellulose ester filter membrane for sterilization due to potential for adsorption.]

ASSAY

• PROCEDURE

Solution A: Add 2.72 g of dibasic potassium phosphate to 800 mL of water. Adjust with phosphoric acid to a pH of 5.8. Add 200 mL of methanol and mix well.

Mobile phase: See [Table 1](#).

Table 1

Time (min)	Methanol (%)	Solution A (%)
0.0	20	80
3.0	20	80
28.0	50	50
28.5	55	45
34.0	55	45
34.5	20	80
40	20	80

Standard solution: 2 mcg/mL of fentanyl prepared from [USP Fentanyl Citrate RS](#) in water

Sample solution: Transfer 1 mL of Injection into a 25-mL volumetric flask, and dilute with water to volume.

Chromatographic system

(See [Chromatography \(621\), System Suitability](#).)

Mode: LC

Detector: UV 210 nm

Column: 4.6-mm \times 10-cm; 2.6- μ m packing L1

Column temperature: 40°

Flow rate: 0.8 mL/min

Injection volume: 20 μ L

System suitability

Sample: Standard solution

[NOTE—The retention time for fentanyl is about 24.6 min.]

Suitability requirements**Tailing factor:** NMT 2.0**Relative standard deviation:** NMT 2.0% for replicate injections**Analysis****Samples:** Standard solution and Sample solutionCalculate the percentage of the labeled amount of fentanyl ($C_{22}H_{28}N_2O$) in the portion of Injection taken:

$$\text{Result} = (r_u/r_s) \times (C_s/C_u) \times 100$$

 r_u = peak response of fentanyl from the Sample solution r_s = peak response of fentanyl from the Standard solution C_s = concentration of fentanyl in the Standard solution (mcg/mL) C_u = nominal concentration of fentanyl in the Sample solution (mcg/mL)**Acceptance criteria:** 90.0%–110.0%**SPECIFIC TESTS**

- **pH (791):** 4.0–7.5
- **STERILITY TESTS (71), Test for Sterility of the Product to Be Examined, Membrane Filtration:** Meets the requirements
- **BACTERIAL ENDOTOXINS TEST (85):** NMT 50.0 USP Endotoxin Units/mg of fentanyl
- **PARTICULATE MATTER IN INJECTIONS (788):** Meets the requirements

ADDITIONAL REQUIREMENTS

- **PACKAGING AND STORAGE:** Package in a light-resistant single-dose container. Store at controlled room temperature or in a refrigerator.

Change to read:

- **Beyond-Use Date:** ▲ In the absence of passing a sterility and endotoxin test, the beyond-use dates in [Pharmaceutical Compounding—Sterile Preparations \(797\)](#) apply. After successful completion of sterility and endotoxin testing, NMT 90 days after the date on which it was compounded when stored at controlled room temperature or in a refrigerator.▲ (CN 1-May-2020)
- **LABELING:** Label it to indicate that it is for use in a single patient only, and to state the Beyond-Use Date.
- **USP REFERENCE STANDARDS (11)**
[USP Fentanyl Citrate RS](#)

Auxiliary Information - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
FENTANYL CITRATE COMPOUNDED INJECTION	Brian Serumaga Science Program Manager	CMP2020 Compounding 2020

Chromatographic Database Information: [Chromatographic Database](#)**Most Recently Appeared In:**

Pharmacopeial Forum: Volume No. 50(1)

Current DocID: GUID-F63BC4D0-6225-4D78-9830-F891DEA4772E_4_en-US**DOI:** https://doi.org/10.31003/USPNF_M10965_04_01**DOI ref:** [ypk9e](#)