

Status: Currently Official on 15-Feb-2025  
 Official Date: Official Prior to 2013  
 Document Type: USP Monographs  
 DocId: GUID-6E9F08E5-CE82-41F4-A8CC-A42D9B36849A\_2\_en-US  
 DOI: [https://doi.org/10.31003/USPNF\\_M34222\\_02\\_01](https://doi.org/10.31003/USPNF_M34222_02_01)  
 DOI Ref: 07zu0

© 2025 USPC  
 Do not distribute

## Fluticasone Propionate Cream

### DEFINITION

Fluticasone Propionate Cream contains NLT 90.0% and NMT 110.0% of the labeled amount of fluticasone propionate ( $C_{25}H_{31}F_3O_5S$ ).

### IDENTIFICATION

- A. [THIN-LAYER CHROMATOGRAPHIC IDENTIFICATION TEST \(201\)](#)

**Standard solution:** 0.4 mg/mL of [USP Fluticasone Propionate RS](#) in acetonitrile

**Test solution:** Transfer a quantity of Cream, equivalent to 1000  $\mu$ g of fluticasone propionate, to a 125-mL separatory funnel. Add 25 mL of acetonitrile and 25 mL of hexane to the separatory funnel. Stopper and shake the funnel until the Cream is completely dispersed. Shake the separatory funnel for an additional 3 min, and allow the phases to separate. Filter the lower layer through a 20-mL syringe containing a cotton plug into a 50-mL volumetric flask. Repeat the extraction with one 7-mL aliquot of acetonitrile, and filter the lower layer into the volumetric flask. Wash the cotton plug with 2 mL of acetonitrile, and collect the washings into the volumetric flask. Dilute the sample extract with acetonitrile to volume. Transfer 12 mL of the sample extract to a glass tube suitable for evaporation, and evaporate to dryness at about 40°. Dissolve the residue in 0.6 mL of acetonitrile. [Note—The *Test solution* may be cloudy because of the presence of undissolved excipients.]

**Chromatographic system**

(See [Chromatography \(621\), Thin-Layer Chromatography](#).)

**Adsorbent:** 0.2-mm layer of chromatographic silica gel mixture on a high-performance thin-layer chromatographic plate, 5- $\mu$ m particle size

**Application volume:** 40  $\mu$ L

**Developing solvent system:** Dichloromethane, ethyl acetate, and glacial acetic acid (30:8:1)

**Analysis**

**Samples:** *Standard solution* and *Test solution*

Separately apply the *Standard solution* and the *Test solution* to the plate. On the same plate, apply 20  $\mu$ L of the *Standard solution*, allow the application to dry, and apply 20  $\mu$ L of the *Test solution* on top of the dried 20- $\mu$ L *Standard solution* spot. Allow each of the applications to dry thoroughly. Place the plate in a tank equilibrated with the developing solvent, and allow the developing solvent to travel about 8 cm from the point of application. Remove the plate and allow to air-dry. Examine the plate under ultraviolet light at 254 nm.

**Acceptance criteria:** The  $R_F$  value of the principal spot from the *Test solution* corresponds to that of the *Standard solution*. [Note—If the excipients in the Cream interfere with the appearance of the principal spot obtained for the *Test solution*, use the *Standard solution* and the *Test solution* overspot to confirm identity.]

- B. The retention time of the major peak of the *Sample solution* corresponds to that of the *Standard solution*, as obtained in the *Assay*.

### ASSAY

- **PROCEDURE**

[Note—Protect the *Standard solution* and the *Sample solution* from direct light by using a light-protective volumetric flask and autosampler vials.]

**Buffer:** 1.2 g/L of monobasic ammonium phosphate. Adjust with phosphoric acid to a pH of 3.50  $\pm$  0.03.

**Mobile phase:** Methanol, acetonitrile, and *Buffer* (46:14:40)

**Diluent:** Alcohol and water (65:35)

**System suitability stock solution:** 0.5 mg/mL of [USP Fluticasone Propionate Nasal Spray Resolution Mixture RS](#) in methanol. [Note—[USP Fluticasone Propionate Nasal Spray Resolution Mixture RS](#) is a mixture of fluticasone propionate and fluticasone propionate related compound D.]

**System suitability solution:** 10  $\mu$ g/mL of [USP Fluticasone Propionate Nasal Spray Resolution Mixture RS](#) in *Diluent* from *System suitability stock solution*

**Standard stock solution:** 0.5 mg/mL [USP Fluticasone Propionate RS](#). Dissolve the standard first in a volume of methanol, equivalent to 80% of the final volume, and dilute with water to volume.

**Standard solution:** 20  $\mu$ g/mL [USP Fluticasone Propionate RS](#) in *Diluent*, from *Standard stock solution*

**Sample solution:** Equivalent to 20 µg/mL of fluticasone propionate. Transfer a quantity of Cream, equivalent to 1000 µg of fluticasone propionate, to a 125-mL separatory funnel. Add to the separatory funnel 25 mL of *Diluent*. Stopper and shake vigorously until the Cream is completely dispersed. Add 25 mL of hexane, shake for an additional 3 min, and allow the phases to separate. Filter the lower layer through a 20-mL syringe containing a cotton plug into a 50-mL volumetric flask. Repeat the extraction with one 5-mL and one 2-mL aliquot of *Diluent*, filtering the lower layers into the volumetric flask. Wash the cotton plug with 1 mL of *Diluent*, and collect the washings into the volumetric flask. Dilute the sample with *Diluent* to volume.

### Chromatographic system

(See [Chromatography \(621\), System Suitability](#).)

**Mode:** LC

**Detector:** 240 nm

**Column:** 4.6-mm × 25-cm; 5-µm packing L1

**Column temperature:** 50°

**Flow rate:** 1.5 mL/min

**Injection size:** 20 µL

### System suitability

**Samples:** System suitability solution and Standard solution

#### Suitability requirements

**Resolution:** NLT 1.4 between fluticasone propionate and fluticasone propionate related compound D, *System suitability solution*

**Tailing factor:** NMT 1.4, *Standard solution* (calculated using the width of the peak at 10% of the height)

**Relative standard deviation:** NMT 2.0%, *Standard solution*

### Analysis

**Samples:** Standard solution and Sample solution

Calculate the percentage of fluticasone propionate ( $C_{25}H_{31}F_3O_5S$ ) in the portion of Cream taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times 100$$

$r_U$  = peak response of fluticasone propionate from the *Sample solution*

$r_S$  = peak response of fluticasone propionate from the *Standard solution*

$C_S$  = concentration of [USP Fluticasone Propionate RS](#) in the *Standard solution* (µg/mL)

$C_U$  = nominal concentration of fluticasone propionate in the *Sample solution* (µg/mL)

**Acceptance criteria:** 90.0%–110.0%

### PERFORMANCE TESTS

- [MINIMUM FILL \(755\)](#): Meets the requirements

### SPECIFIC TESTS

- [pH \(791\)](#): 4.5–6.5
- [MICROBIAL ENUMERATION TESTS \(61\)](#) and [TESTS FOR SPECIFIED MICROORGANISMS \(62\)](#): Meets the requirements of the tests for absence of *Staphylococcus aureus* and *Pseudomonas aeruginosa*. The total aerobic microbial count is NMT 100 cfu/g, and the total combined molds and yeasts count is NMT 10 cfu/g.

### ADDITIONAL REQUIREMENTS

- **PACKAGING AND STORAGE:** Preserve in collapsible tubes or tight containers, protected from light. Store between 2° and 30°.
- [USP REFERENCE STANDARDS \(11\)](#).
  - [USP Fluticasone Propionate RS](#)
  - [USP Fluticasone Propionate Nasal Spray Resolution Mixture RS](#)

This Reference Standard is a mixture of fluticasone propionate and fluticasone propionate related compound D, and the chemical names for both are given below:

*Fluticasone propionate*: S-Fluoromethyl 6 $\alpha$ ,9 $\alpha$ -difluoro-11 $\beta$ -hydroxy-16 $\alpha$ -methyl-3-oxo-17 $\alpha$ -propionyloxyandrosta-1,4-diene-17 $\beta$ -carbothioate.

*Fluticasone propionate related compound D*: S-Methyl-6 $\alpha$ ,9 $\alpha$ -difluoro-11 $\beta$ -hydroxy-16 $\alpha$ -methyl-3-oxo-17 $\alpha$ -propionyloxy-androsta-1,4-diene-17 $\beta$ -carbothioate.

Topic/Question	Contact	Expert Committee
FLUTICASONE PROPIONATE CREAM	<a href="#">Documentary Standards Support</a>	SM52020 Small Molecules 5

**Chromatographic Database Information:** [Chromatographic Database](#)

**Most Recently Appeared In:**

Pharmacopeial Forum: Volume No. 45(6)

**Current DocID: GUID-6E9F08E5-CE82-41F4-A8CC-A42D9B36849A\_2\_en-US**

**Previous DocID: GUID-6E9F08E5-CE82-41F4-A8CC-A42D9B36849A\_1\_en-US**

**DOI: [https://doi.org/10.31003/USPNF\\_M34222\\_02\\_01](https://doi.org/10.31003/USPNF_M34222_02_01)**

**DOI ref: 07zu0**

OFFICIAL