

Status: Currently Official on 15-Feb-2025
Official Date: Official Prior to 2013
Document Type: Reagents
DocId: GUID-7734C358-4409-42F2-8D70-A862404F1A64_1_en-US
DOI: https://doi.org/10.31003/USPNF_R3174_01_01
DOI Ref: lg8pb

© 2025 USPC
Do not distribute

Fuchsin–Pyrogallol TS

–Dissolve 100 mg of basic fuchsin in 50 mL of water that previously has been boiled for 15 minutes and allowed to cool slightly. Cool, add 2 mL of a saturated solution of sodium bisulfite, mix, and allow to stand for not less than 3 hours. Add 0.9 mL of hydrochloric acid, mix, and allow to stand overnight. Add 100 mg of pyrogallol, shake until solution is effected, and dilute with water to 100 mL. Store in an amber-colored glass bottle in a refrigerator.

Auxiliary Information - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
FUCHSIN–PYROGALLOL TS	Margareth R.C. Marques Principal Scientific Liaison	HDQ Headquarters

Most Recently Appeared In:

Pharmacopeial Forum: Volume No. Information currently unavailable

Current DocID: [GUID-7734C358-4409-42F2-8D70-A862404F1A64_1_en-US](#)

DOI: https://doi.org/10.31003/USPNF_R3174_01_01

DOI ref: [lg8pb](#)

OFFICIAL