

Status: Currently Official on 15-Feb-2025  
Official Date: Official Prior to 2013  
Document Type: Reagents  
DocId: GUID-3ECE90F7-D964-4572-8255-7C756C24948B\_1\_en-US  
DOI: [https://doi.org/10.31003/USPNF\\_R3175\\_01\\_01](https://doi.org/10.31003/USPNF_R3175_01_01)  
DOI Ref: qf7av

© 2025 USPC  
Do not distribute

## Fuchsin–Sulfurous Acid TS

–Dissolve 200 mg of basic fuchsin in 120 mL of hot water, and allow the solution to cool. Add a solution of 2 g of anhydrous sodium sulfite in 20 mL of water, then add 2 mL of hydrochloric acid. Dilute the solution with water to 200 mL, and allow to stand for at least 1 hour. Prepare this solution fresh.

**Auxiliary Information** - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
FUCHSIN–SULFUROUS ACID TS	<a href="#">Margareth R.C. Marques</a> Principal Scientific Liaison	HDQ Headquarters

**Most Recently Appeared In:**

Pharmacopeial Forum: Volume No. Information currently unavailable

**Current DocId:** [GUID-3ECE90F7-D964-4572-8255-7C756C24948B\\_1\\_en-US](#)

**DOI:** [https://doi.org/10.31003/USPNF\\_R3175\\_01\\_01](https://doi.org/10.31003/USPNF_R3175_01_01)

**DOI ref:** [qf7av](#)

OFFICIAL