

Status: Currently Official on 15-Feb-2025  
Official Date: Official Prior to 2013  
Document Type: USP Monographs  
DocId: GUID-C393351E-17A3-4289-BE2D-A42D13040AE1\_1\_en-US  
DOI: [https://doi.org/10.31003/USPNF\\_M38470\\_01\\_01](https://doi.org/10.31003/USPNF_M38470_01_01)  
DOI Ref: o37bt

© 2025 USPC  
Do not distribute

## Hydrocortisone Valerate Cream

### DEFINITION

Hydrocortisone Valerate Cream is Hydrocortisone Valerate in a suitable cream base. It contains NLT 90.0% and NMT 110.0% of the labeled amount of hydrocortisone valerate ( $C_{26}H_{38}O_6$ ).

### IDENTIFICATION

- **A. THIN-LAYER CHROMATOGRAPHIC IDENTIFICATION TEST (201).**

**Samples:** Standard solution and Sample solution from the Assay

**Acceptance criteria:** Meets the requirements

### ASSAY

- **PROCEDURE**

**Mobile phase:** Acetonitrile and water (45:55)

**Internal standard solution:** 2.0 mg/mL of ethyl benzoate in methanol

**Standard stock solution:** 0.5 mg/mL of [USP Hydrocortisone Valerate RS](#) in methanol. Prepare immediately before use.

**Standard solution:** 0.1 mg/mL of [USP Hydrocortisone Valerate RS](#) in methanol, prepared as follows. Pipet 2 mL of the Standard stock solution and 2 mL of the Internal standard solution into a 10-mL volumetric flask, and dilute with methanol to volume.

**Sample solution:** Nominally 0.1 mg/mL of hydrocortisone valerate, prepared as follows. Transfer a quantity of Cream, equivalent to 1 mg of hydrocortisone valerate, to a screw-capped tube. Add 8.0 mL of a mixture of methanol and water (3:1), and swirl to disperse. Heat at 80° for 1 min, swirl again, and allow to cool to room temperature. Add 2.0 mL of the Internal standard solution. Centrifuge for 5 min, and filter, if necessary, to obtain a clear supernatant.

#### Chromatographic system

(See [Chromatography \(621\), System Suitability](#).)

**Mode:** LC

**Detector:** UV 254 nm

**Column:** 3.9-mm × 30-cm; packing L1

**Flow rate:** 1 mL/min

**Injection volume:** 10  $\mu$ L

#### System suitability

**Sample:** Standard solution

[NOTE—The relative retention times for ethyl benzoate and hydrocortisone valerate are about 0.8 and 1.0, respectively.]

#### Suitability requirements

**Resolution:** NLT 3.0 between ethyl benzoate and hydrocortisone valerate

**Relative standard deviation:** NMT 2.0%

#### Analysis

**Samples:** Standard solution and Sample solution

Calculate the percentage of the labeled amount of hydrocortisone valerate ( $C_{26}H_{38}O_6$ ) in the portion of Cream taken:

$$\text{Result} = (R_U/R_S) \times (C_S/C_U) \times 100$$

$R_U$  = peak response ratio of hydrocortisone valerate to the internal standard from the Sample solution

$R_S$  = peak response ratio of hydrocortisone valerate to the internal standard from the Standard solution

$C_S$  = concentration of [USP Hydrocortisone Valerate RS](#) in the Standard solution (mg/mL)

$C_U$  = nominal concentration of hydrocortisone valerate in the Sample solution (mg/mL)

#### PERFORMANCE TESTS

- MINIMUM FILL (755): Meets the requirements

#### ADDITIONAL REQUIREMENTS

- **PACKAGING AND STORAGE**: Preserve in well-closed containers.
- USP REFERENCE STANDARDS (11)  
[USP Hydrocortisone Valerate RS](#)

**Auxiliary Information** - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
HYDROCORTISONE VALERATE CREAM	<a href="#">Documentary Standards Support</a>	SM52020 Small Molecules 5

**Chromatographic Database Information:** [Chromatographic Database](#)

#### Most Recently Appeared In:

Pharmacopeial Forum: Volume No. Information currently unavailable

**Current DocID: GUID-C393351E-17A3-4289-BE2D-A42D13040AE1\_1\_en-US**

**DOI:** [https://doi.org/10.31003/USPNF\\_M38470\\_01\\_01](https://doi.org/10.31003/USPNF_M38470_01_01)

**DOI ref:** o37bt

OFFICIAL