

Status: Currently Official on 14-Feb-2025
 Official Date: Official as of 27-Mar-2024
 Document Type: USP Monographs
 DocId: GUID-98FFD64B-5EBE-4474-A556-E1996987BA0E_2_en-US
 DOI: https://doi.org/10.31003/USPNF_M39430_02_01
 DOI Ref: 9ueaz

© 2025 USPC
 Do not distribute

Hydroxyurea Capsules

To view the Notice from the Expert Committee that posted in conjunction with this accelerated revision, please click
<https://www.uspnf.com/rb-hydroxyurea-caps-20240426>.

DEFINITION

Hydroxyurea Capsules contain NLT 90.0% and NMT 110.0% of the labeled amount of hydroxyurea ($\text{CH}_4\text{N}_2\text{O}_2$).

IDENTIFICATION

- A.

Standard: Transfer 30 mg of [USP Hydroxyurea RS](#) to a suitable centrifuge tube, and add 10 mL of anhydrous [methanol](#). Centrifuge for 3 min. Transfer 1.0 mL of the clear supernatant to a mortar containing 500 mg of [potassium bromide](#), triturate to a homogeneous blend, dry in a vacuum desiccator at 60° for 3 h, and prepare a suitable disk.

Sample: Transfer a portion of Capsule contents, equivalent to 30 mg of hydroxyurea, to a suitable centrifuge tube, and add 10 mL of anhydrous [methanol](#). Centrifuge for 3 min. Transfer 1.0 mL of the clear supernatant to a mortar containing 500 mg of [potassium bromide](#), triturate to a homogeneous blend, dry in a vacuum desiccator at 60° for 3 h, and prepare a suitable disk.

Acceptance criteria: The IR absorption spectrum of the *Sample* exhibits maxima only at the same wavenumbers as that of the *Standard*.

ASSAY

- PROCEDURE

Solution A: Dissolve 1.7 g of [tetrabutylammonium hydrogen sulfate](#) and 1.74 g of [dibasic potassium phosphate](#) in 1000 mL of [water](#). Adjust with 1 N [sodium hydroxide](#) or [phosphoric acid](#) to a pH of 5.0.

Mobile phase: [Methanol](#) and Solution A (15:85)

System suitability solution: 0.4 mg/mL each of [USP Hydroxyurea RS](#) and [hydroxylamine hydrochloride](#) in *Mobile phase*

Standard solution: 0.4 mg/mL of [USP Hydroxyurea RS](#) in *Mobile phase*

Sample solution: Nominally 0.4 mg/mL of hydroxyurea in *Mobile phase* prepared as follows. Remove, as completely as possible, the contents of NLT 20 Capsules and grind to a fine powder. Transfer a portion of the powder, equivalent to 200 mg of hydroxyurea, to a 500-mL volumetric flask. Add 300 mL of *Mobile phase*, sonicate for 10 min, stir with the aid of a magnetic stirrer for 30 min, sonicate for an additional 10 min, and dilute with *Mobile phase* to volume. Filter a portion of the resulting solution, discarding the first 2 mL of the filtrate.

Chromatographic system

(See [Chromatography \(621\), System Suitability](#).)

Mode: LC

Detector: UV 214 nm

Column: 4.6-mm × 25-cm; 5-μm packing [L1](#)

Flow rate: 0.5 mL/min

Injection volume: 10 μL

System suitability

Samples: *System suitability solution* and *Standard solution*

Suitability requirements

Resolution: NLT 1.5 between hydroxylamine and hydroxyurea, *System suitability solution*

Column efficiency: NLT 5000 for hydroxyurea, *System suitability solution*

Tailing factor: NMT 1.5 for hydroxyurea, *System suitability solution*

Relative standard deviation: NMT 2.0%, *Standard solution*

Analysis

Samples: *Standard solution* and *Sample solution*

Calculate the percentage of the labeled amount of hydroxyurea ($\text{CH}_4\text{N}_2\text{O}_2$) in the portion of Capsules taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times 100$$

r_U = peak response of hydroxyurea from the *Sample solution*

r_S = peak response of hydroxyurea from the *Standard solution*

C_S = concentration of [USP Hydroxyurea RS](#) in the *Standard solution* (mg/mL)

C_U = nominal concentration of hydroxyurea in the *Sample solution* (mg/mL)

Acceptance criteria: 90.0%–110.0%

PERFORMANCE TESTS

Change to read:

- [Dissolution \(711\)](#)

▲ Test 1 ▲ (RB 27-Mar-2024)

Medium: [Water](#); 500 mL

Apparatus 2: 50 rpm

Time: 30 min

Analysis: Calculate the percentage of the labeled amount of hydroxyurea ($\text{CH}_4\text{N}_2\text{O}_2$) dissolved by using the procedure set forth in the Assay, making any necessary modifications.

Tolerances: NLT 80% (Q) of the labeled amount of hydroxyurea ($\text{CH}_4\text{N}_2\text{O}_2$) is dissolved.

▲ Test 2: If the product complies with this test, the labeling indicates that it meets USP *Dissolution Test 2*.

Medium: 0.1 N [hydrochloric acid](#); 500 mL

Apparatus 2: 50 rpm

Time: 30 min

Buffer: Dissolve 1.7 g of [tetrabutylammonium hydrogen sulfate](#) and 1.74 g of [dibasic potassium phosphate](#) in 1000 mL of [water](#). Adjust with [phosphoric acid](#) to a pH of 5.0.

Mobile phase: [Methanol](#) and **Buffer** (15:85)

Standard solution: ($L/500$) mg/mL of [USP Hydroxyurea RS](#) in **Medium**, where L is the label claim in mg/Capsule

Sample solution: Pass a portion of the solution under test through a suitable filter of 0.45- μm pore size, discarding an appropriate volume of filtrate so that a consistent result can be obtained.

Chromatographic system

(See [Chromatography \(621\), System Suitability](#).)

Mode: LC

Detector: UV 214 nm

Column: 4.6-mm \times 25-cm; 5- μm packing [L1](#)

Column temperature: 30°

Flow rate: 0.5 mL/min

Injection volume: 10 μL

Run time: NLT 4.5 times the retention time of hydroxyurea

System suitability

Sample: *Standard solution*

Suitability requirements

Relative standard deviation: NMT 2.0%

Analysis

Samples: *Standard solution* and *Sample solution*

Calculate the percentage of the labeled amount of hydroxyurea ($\text{CH}_4\text{N}_2\text{O}_2$) dissolved:

$$\text{Result} = (r_U/r_S) \times C_S \times V \times (1/L) \times 100$$

r_U = peak response of hydroxyurea from the *Sample solution*

r_S = peak response of hydroxyurea from the *Standard solution*

C_S = concentration of [USP Hydroxyurea RS](#) in the *Standard solution* (mg/mL)

V = volume of **Medium**, 500 mL

L = label claim (mg/Capsule)

Tolerances: NLT 80% (Q) of the labeled amount of hydroxyurea ($\text{CH}_4\text{N}_2\text{O}_2$) is dissolved.▲ (RB 27-Mar-2024)

- [Uniformity of Dosage Units \(905\)](#): Meet the requirements

ADDITIONAL REQUIREMENTS

- **PACKAGING AND STORAGE:** Preserve in tight containers, in a dry atmosphere.

Add the following:

- ▲ **LABELING:** The labeling states the *Dissolution* test used only if *Test 1* is not used.▲ (RB 27-Mar-2024)

- [USP Reference Standards \(11\)](#)

[USP Hydroxyurea RS](#)

Auxiliary Information - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
HYDROXYUREA CAPSULES	Documentary Standards Support	SM32020 Small Molecules 3

Chromatographic Database Information: [Chromatographic Database](#)

Most Recently Appeared In:

Pharmacopeial Forum: Volume No. PF 29(6)

Current DocID: GUID-98FFD64B-5EBE-4474-A556-E1996987BA0E_2_en-US

DOI: https://doi.org/10.31003/USPNF_M39430_02_01

DOI ref: 9ueaz

OFFICIAL