

Status: Currently Official on 15-Feb-2025
Official Date: Official Prior to 2013
Document Type: USP Monographs
DocId: GUID-A95105D9-7BB9-42D7-BAE0-69B49D51F0F5_1_en-US
DOI: https://doi.org/10.31003/USPNF_M39710_01_01
DOI Ref: 7pc9s

© 2025 USPC
Do not distribute

Hyoscyamine Sulfate Oral Solution

DEFINITION

Hyoscyamine Sulfate Oral Solution contains NLT 90.0% and NMT 110.0% of the labeled amount of hyoscyamine sulfate $[(C_{17}H_{23}NO_3)_2 \cdot H_2SO_4 \cdot 2H_2O]$.

IDENTIFICATION

- **A.** The retention time of the major peak of the *Sample solution* corresponds to that of the *Standard solution*, as obtained in the Assay.

ASSAY

• PROCEDURE

Buffer: Transfer 13.6 g of monobasic potassium phosphate to a 2000-mL volumetric flask, dissolve in about 1800 mL of water, adjust with phosphoric acid to a pH of 3.0 ± 0.1 , dilute with water to volume, and filter.

Mobile phase: With continuous stirring, add 0.3 mL of triethylamine to 1800 mL of *Buffer*. Add 200 mL of acetonitrile, and degas.

Diluent: 0.01 N hydrochloric acid

Standard stock solution: 0.16 mg/mL of anhydrous hyoscyamine sulfate from [USP Hyoscyamine Sulfate RS](#) in *Diluent*. [NOTE—This solution may be stored in a refrigerator for 30 days.]

Standard solution: 4.8 μ g/mL of anhydrous hyoscyamine sulfate from the *Standard stock solution* in *Diluent*

Sample solution: 5 μ g/mL of hyoscyamine sulfate from an appropriate volume of Oral Solution in *Diluent*. Pass an aliquot through a suitable filter of 0.45- μ m pore size, and discard the first 5 mL of the filtrate.

Chromatographic system

(See [Chromatography \(621\), System Suitability](#).)

Mode: LC

Detector: UV 205 nm

Columns

Guard: 3-mm \times 4-mm; packing L11

Analytical: 4.6-mm \times 15-cm; 4- μ m packing L11

Column temperature: 30°

Flow rate: 1.0 mL/min

Injection volume: 50 μ L

System suitability

Sample: *Standard solution*

Suitability requirements

Tailing factor: NMT 1.8

Relative standard deviation: NMT 2.0% for six replicate injections

Analysis

Samples: *Standard solution* and *Sample solution*

Calculate the percentage of the labeled amount of hyoscyamine sulfate $[(C_{17}H_{23}NO_3)_2 \cdot H_2SO_4 \cdot 2H_2O]$ in the portion of Oral Solution taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times (M_{r1}/M_{r2}) \times 100$$

r_U = peak response from the *Sample solution*

r_S = peak response from the *Standard solution*

C_S = concentration of anhydrous hyoscyamine sulfate in the *Standard solution* (μ g/mL)

C_U = nominal concentration of hyoscyamine sulfate in the *Sample solution* (μ g/mL)

M_{r1} = molecular weight of hyoscyamine sulfate, 712.85

M_{r2} = molecular weight of anhydrous hyoscyamine sulfate, 676.83

Acceptance criteria: 90.0%–110.0%

PERFORMANCE TESTS

- [UNIFORMITY OF DOSAGE UNITS \(905\)](#)

For single-unit containers

Acceptance criteria: Meets the requirements

- [DELIVERABLE VOLUME \(698\)](#)

For multiple-unit containers

Acceptance criteria: Meets the requirements

SPECIFIC TESTS

- [pH \(791\)](#): 3.0–6.5

ADDITIONAL REQUIREMENTS

- **PACKAGING AND STORAGE:** Preserve in tight, light-resistant containers, and store at controlled room temperature.

- [USP REFERENCE STANDARDS \(11\)](#)

[USP Hyoscyamine Sulfate RS](#)

Auxiliary Information - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
HYOSCYAMINE SULFATE ORAL SOLUTION	<u>Documentary Standards Support</u>	SM32020 Small Molecules 3
REFERENCE STANDARD SUPPORT	RS Technical Services <u>RSTECH@usp.org</u>	SM32020 Small Molecules 3

Chromatographic Database Information: [Chromatographic Database](#)

Most Recently Appeared In:

Pharmacopeial Forum: Volume No. PF 31(5)

Current DocID: GUID-A95105D9-7BB9-42D7-BAE0-69B49D51F0F5_1_en-US

DOI: https://doi.org/10.31003/USPNF_M39710_01_01

DOI ref: [7pc9s](#)