

Status: Currently Official on 15-Feb-2025
Official Date: Official Prior to 2013
Document Type: Reagents
DocId: GUID-12BB983F-2857-4A2E-B121-4B2D6F585166_1_en-US
DOI: https://doi.org/10.31003/USPNF_R3187_01_01
DOI Ref: 23lu3

© 2025 USPC
Do not distribute

Indophenol–Acetate TS

(for the assay of *Corticotropin Injection*)—To 60 mL of standard dichlorophenol-indophenol solution (see in the section *Volumetric Solutions*) add water to make 250 mL. Add to the resulting solution an equal volume of sodium acetate solution freshly prepared by dissolving 13.66 g of anhydrous sodium acetate in water to make 500 mL and adjusting with 0.5 N acetic acid to a pH of 7. Store in a refrigerator, and use within 2 weeks.

Auxiliary Information - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
INDOPHENOL–ACETATE TS	Margareth R.C. Marques Principal Scientific Liaison	HDQ Headquarters

Most Recently Appeared In:

Pharmacopeial Forum: Volume No. Information currently unavailable

Current DocID: [GUID-12BB983F-2857-4A2E-B121-4B2D6F585166_1_en-US](#)

DOI: https://doi.org/10.31003/USPNF_R3187_01_01

DOI ref: [23lu3](#)

OFFICIAL