

Status: Currently Official on 15-Feb-2025
Official Date: Official Prior to 2013
Document Type: Reagents
DocId: GUID-190BB90A-8350-41D4-82E0-88920705A993_1_en-US
DOI: https://doi.org/10.31003/USPNF_R3188_01_01
DOI Ref: e6rsi

© 2025 USPC
Do not distribute

Intestinal Fluid, Simulated, TS

—Dissolve 6.8 g of monobasic potassium phosphate in 250 mL of water, mix, and add 77 mL of 0.2 N sodium hydroxide and 500 mL of water. Add 10.0 g of pancreatin, mix, and adjust the resulting solution with either 0.2 N sodium hydroxide or 0.2 N hydrochloric acid to a pH of 6.8 ± 0.1 . Dilute with water to 1000 mL.

Auxiliary Information - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
INTESTINAL FLUID, SIMULATED, TS	Margareth R.C. Marques Principal Scientific Liaison	HDQ Headquarters

Most Recently Appeared In:

Pharmacopeial Forum: Volume No. Information currently unavailable

Current DocId: [GUID-190BB90A-8350-41D4-82E0-88920705A993_1_en-US](#)

DOI: https://doi.org/10.31003/USPNF_R3188_01_01

DOI ref: [e6rsi](#)

OFFICIAL