

Status: Currently Official on 15-Feb-2025
Official Date: Official Prior to 2013
Document Type: Reagents
DocId: GUID-29F88E74-5185-4790-81DF-5D60258A6F5B_1_en-US
DOI: https://doi.org/10.31003/USPNF_R3196_01_01
DOI Ref: I8fdy

© 2025 USPC
Do not distribute

Iron Salicylate TS

—Dissolve 500 mg of ferric ammonium sulfate in 250 mL of water containing 10 mL of diluted sulfuric acid, and add water to make 500 mL. To 100 mL of the resulting solution add 50 mL of a 1.15% solution of sodium salicylate, 20 mL of diluted acetic acid, and 80 mL of a 13.6% solution of sodium acetate, then add water to make 500 mL. Store in a well-closed container. Protect from light. Use within 2 weeks.

Auxiliary Information - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
IRON SALICYLATE TS	Margareth R.C. Marques Principal Scientific Liaison	HDQ Headquarters

Most Recently Appeared In:

Pharmacopeial Forum: Volume No. Information currently unavailable

Current DocID: [GUID-29F88E74-5185-4790-81DF-5D60258A6F5B_1_en-US](#)

DOI: https://doi.org/10.31003/USPNF_R3196_01_01

DOI ref: [I8fdy](#)

