

Status: Currently Official on 15-Feb-2025
Official Date: Official Prior to 2013
Document Type: USP Monographs
DocId: GUID-B93FE304-8646-4454-98CC-FE212D966CD5_1_en-US
DOI: https://doi.org/10.31003/USPNF_M43738_01_01
DOI Ref: wa5qq

© 2025 USPC
Do not distribute

Ivermectin Topical Solution

» Ivermectin Topical Solution is a topical solution in a suitable vehicle. It contains not less than 95.0 percent and not more than 105.0 percent of the labeled amount of Ivermectin, calculated as the sum of component H_2B_{1a} ($C_{48}H_{74}O_{14}$) plus component H_2B_{1b} ($C_{47}H_{72}O_{14}$). The ratio of the contents, $H_2B_{1a}/(H_2B_{1a} + H_2B_{1b})$, is not less than 90.0 percent.

Packaging and storage—Preserve in well-closed containers. Store at a temperature not more than 30°.

Labeling—Label it to indicate that it is for veterinary use only.

USP REFERENCE STANDARDS (11)—

[USP Ivermectin RS](#)

Identification—

A: *Thin-Layer Chromatographic Identification Test (201)*—

Test solution—Dissolve a volume of Topical Solution in methanol, and dilute quantitatively, and stepwise if necessary, with methanol to obtain a solution containing 0.5 mg per mL of ivermectin, and filter.

Injection volume: 2 μ L.

Developing solvent system: unsaturated chamber, freshly prepared and equilibrated mixture of methylene chloride, methanol, and ammonium hydroxide (90:9:1).

Procedure—Remove the plate, allow to air dry, and examine under short- and long-wavelength UV light: the retardation factor, R_F , of the principal spot obtained from the *Test solution* corresponds to that obtained from the *Standard solution*.

B: The retention times of the two principal component peaks of ivermectin in the chromatogram of the *Assay preparation* corresponds to that of the two principal component peaks of ivermectin in the chromatogram of the *Standard preparation*, obtained as directed in the *Assay*.

Chromatographic purity—

Mobile phase and Chromatographic system—Proceed as directed in the *Assay*.

Standard solution—Dissolve an accurately weighed quantity of [USP Ivermectin RS](#) in methanol, and dilute quantitatively, and stepwise if necessary, with methanol to obtain a 0.004 mg per mL solution.

Test solution—Dissolve a quantity of Topical Solution in methanol, and dilute quantitatively, and stepwise if necessary, with methanol to obtain a solution containing 0.4 mg of ivermectin per mL of solution, based on the label claim.

Procedure—Inject equal volumes (about 20 μ L) of the *Test solution* and the *Standard solution* into the chromatograph, record the chromatograms, and measure all of the peak responses. Calculate the percentage of each impurity in the portion of Topical Solution taken by the formula:

$$100(C_s/C_T) (r_i/r_s)$$

in which C_s is the concentration, in mg per mL, of ivermectin in the *Standard solution*; C_T is the concentration, in mg per mL, of ivermectin in the *Test solution*; r_i is the peak response for each impurity from the *Test solution*; and r_s is the ivermectin peak response obtained from *Standard solution*. Not more than 2.7% of the peak with a relative retention time of about 1.3 to 1.5 to that of the principal peak is found; not more than 1.0% of any other impurity is found; and not more than 6.0% of total impurities is found. Disregard any peak below 0.05%.

Assay—

Mobile phase—Prepare a filtered and degassed mixture of acetonitrile, methanol, and water (106:55:39). Make adjustments if necessary (see *System Suitability* under [Chromatography \(621\)](#)).

Standard preparation—Dissolve an accurately weighed quantity of [USP Ivermectin RS](#) in methanol, and dilute quantitatively, and stepwise if necessary, with methanol to obtain a solution having a known concentration of about 0.4 mg per mL.

Assay preparation—Dilute a volume of Topical Solution, quantitatively, and stepwise if necessary, with methanol to obtain a solution containing 0.4 mg of ivermectin per mL of solution, based on the label claim.

Chromatographic system (see [CHROMATOGRAPHY \(621\)](#))—The liquid chromatograph is equipped with a 245-nm detector and a 4.6-mm \times 25-cm column that contains packing L1. The flow rate is about 1.5 mL per minute. Chromatograph the *Standard preparation*, and record the peak

responses as directed for *Procedure*: the resolution, R , between the first peak (component H_2B_{1b}) and the second peak (component H_2B_{1a}) is not less than 3.0; and the relative standard deviation for replicate injections is not more than 2.0%, determined from the H_2B_{1a} peak.

Procedure—Separately inject equal volumes (about 20 μ L) of the *Standard preparation* and the *Assay preparation* into the chromatograph, record the chromatograms, and measure the responses for the component H_2B_{1a} plus component H_2B_{1b} . Calculate the percentage of label claim of ivermectin ($H_2B_{1a} + H_2B_{1b}$) in the portion of Topical Solution taken by the formula:

$$100(C_s/C_u)(r_u/r_s)$$

in which C_s is the concentration, in mg per mL, of [USP Ivermectin RS](#) in the *Standard preparation*; C_u is the concentration, in mg per mL, of ivermectin in the *Assay preparation*; r_u is the total peak response of H_2B_{1a} plus H_2B_{1b} peaks obtained from the *Assay preparation*; and r_s is the total peak response of H_2B_{1a} plus H_2B_{1b} peaks obtained from the *Standard preparation*. Calculate the ratio of the contents, in percent, of $H_2B_{1a}/(H_2B_{1a} + H_2B_{1b})$ in the portion of Topical Solution taken by the formula:

$$100(r_1/r_u)$$

in which r_1 is the peak response of H_2B_{1a} obtained from the *Assay preparation*; and r_u is as described above.

Auxiliary Information - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
IVERMECTIN TOPICAL SOLUTION	Documentary Standards Support	SM32020 Small Molecules 3
REFERENCE STANDARD SUPPORT	RS Technical Services RSTECH@usp.org	SM32020 Small Molecules 3

Chromatographic Database Information: [Chromatographic Database](#)

Most Recently Appeared In:

Pharmacopeial Forum: Volume No. PF 33(5)

Current DocID: GUID-B93FE304-8646-4454-98CC-FE212D966CD5_1_en-US

DOI: https://doi.org/10.31003/USPNF_M43738_01_01

DOI ref: [wa5qq](#)