

Status: Currently Official on 15-Feb-2025
Official Date: Official Prior to 2013
Document Type: Reagents
DocId: GUID-28DEA5BE-8FB8-4D1A-BFC3-B6EEA75F35E4_1_en-US
DOI: https://doi.org/10.31003/USPNF_R2147_01_01
DOI Ref: 3l8bh

© 2025 USPC
Do not distribute

Magnesium Sulfate, Anhydrous,

MgSO₄ 120.37 CAS RN[®]: 7487-88-9.—Anhydrous Magnesium Sulfate may be prepared as follows. Place a suitable quantity of magnesium sulfate (see above), preferably powdered, in a shallow vessel, and expose to a temperature of about 80° for several hours with occasional stirring. Then heat at 275° to 300° until the weight is practically constant. Transfer the product while still warm to tight containers, as the anhydrous salt is very hygroscopic.

Auxiliary Information - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
MAGNESIUM SULFATE, ANHYDROUS	Margareth R.C. Marques Principal Scientific Liaison	HDQ Headquarters

Most Recently Appeared In:

Pharmacopeial Forum: Volume No. Information currently unavailable

Current DocId: [GUID-28DEA5BE-8FB8-4D1A-BFC3-B6EEA75F35E4_1_en-US](#)

DOI: https://doi.org/10.31003/USPNF_R2147_01_01

DOI ref: [3l8bh](#)

OFFICIAL