

Status: Currently Official on 16-Feb-2025  
Official Date: Official Prior to 2013  
Document Type: Reagents  
DocId: GUID-C7105857-34B0-460B-B79D-5B1FF7D3E5A1\_1\_en-US  
DOI: [https://doi.org/10.31003/USPNF\\_R3213\\_01\\_01](https://doi.org/10.31003/USPNF_R3213_01_01)  
DOI Ref: swo3y

© 2025 USPC  
Do not distribute

## Mercuric Iodide TS (*Valser's Reagent*)—

Slowly add potassium iodide solution (1 in 10) to red mercuric iodide until almost all of the latter is dissolved, and filter off the excess. A solution containing 10 g of potassium iodide in 100 mL dissolves approximately 14 g of HgI<sub>2</sub> at 20°.

**Auxiliary Information** - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
MERCURIC IODIDE TS (VALSER'S REAGENT)	<a href="#">Margareth R.C. Marques</a> Principal Scientific Liaison	HDQ Headquarters

**Most Recently Appeared In:**

Pharmacopeial Forum: Volume No. Information currently unavailable

**Current DocID:** [GUID-C7105857-34B0-460B-B79D-5B1FF7D3E5A1\\_1\\_en-US](#)

**DOI:** [https://doi.org/10.31003/USPNF\\_R3213\\_01\\_01](https://doi.org/10.31003/USPNF_R3213_01_01)

**DOI ref:** [swo3y](#)

OFFICIAL