


Status: Currently Official on 15-Feb-2025
Official Date: Official as of 01-May-2020
Document Type: USP Monographs
DocId: GUID-91439311-E1CE-4101-A135-52A0D5DC89E1_4_en-US
DOI: https://doi.org/10.31003/USPNF_M51200_04_01
DOI Ref: 74000

© 2025 USPC
Do not distribute

Methocarbamol



C₁₁H₁₅NO₅ 241.24
1,2-Propanediol, 3-(2-methoxyphenoxy)-, 1-carbamate, (±)-;
(±)-3-(o-Methoxyphenoxy)-1,2-propanediol 1-carbamate CAS RN®: 532-03-6; UNII: 125OD7737X.

DEFINITION
Methocarbamol contains NLT 98.5% and NMT 101.5% of methocarbamol (C₁₁H₁₅NO₅), calculated on the dried basis.

IDENTIFICATION
Change to read:
• **A.** ▲ [SPECTROSCOPIC IDENTIFICATION TESTS \(197\)](#), [Infrared Spectroscopy: 197K](#) ▲ (CN 1-MAY-2020)
• **B.** The retention time of the major peak of the *Sample solution* corresponds to that of the *Standard solution*, as obtained in the Assay.

ASSAY
• **PROCEDURE**
Buffer: 6.8 g/L of [monobasic potassium phosphate](#) in water. Adjust with [phosphoric acid](#) or [sodium hydroxide](#) to a pH of 4.5.
Mobile phase: Methanol and *Buffer* (30:70)
System suitability solution: 1.0 mg/mL of [USP Methocarbamol RS](#) and 0.005 mg/mL of [USP Guaifenesin RS](#) in *Mobile phase*
Standard solution: 0.1 mg/mL of [USP Methocarbamol RS](#) in *Mobile phase*
Sample solution: 0.1 mg/mL of Methocarbamol in *Mobile phase*
Chromatographic system
(See [Chromatography \(621\)](#), [System Suitability](#).)
Mode: LC
Detector: UV 274 nm
Column: 4.6-mm × 15-cm; 3-µm packing L1
Column temperature: 30°
Flow rate: 0.8 mL/min
Injection volume: 20 µL
Run time: 1.5 times the retention time of methocarbamol
System suitability
Samples: *System suitability solution* and *Standard solution*
[NOTE—See [Table 1](#) for relative retention times.]
Suitability requirements
Resolution: NLT 3.5 between methocarbamol and guaifenesin, *System suitability solution*
Tailing factor: NMT 2.0, *Standard solution*
Relative standard deviation: NMT 0.73%, *Standard solution*

Analysis
Samples: *Standard solution* and *Sample solution*
Calculate the percentage of methocarbamol (C₁₁H₁₅NO₅) in the portion of Methocarbamol taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times 100$$

r_U = peak response of methocarbamol from the *Sample solution*

r_S = peak response of methocarbamol from the *Standard solution*

C_S = concentration of [USP Methocarbamol RS](#) in the *Standard solution* (mg/mL)

C_U = concentration of Methocarbamol in the *Sample solution* (mg/mL)

Acceptance criteria: 98.5%–101.5% on the dried basis

IMPURITIES

- **RESIDUE ON IGNITION (281):** NMT 0.1%
- **ORGANIC IMPURITIES**

Mobile phase, System suitability solution, and Chromatographic system: Proceed as directed in the Assay.

Standard solution: 0.005 mg/mL of [USP Methocarbamol RS](#) in *Mobile phase*

Sample solution: 1 mg/mL of Methocarbamol in *Mobile phase*

System suitability

Samples: *System suitability solution* and *Standard solution*

[NOTE—See [Table 1](#) for the relative retention times.]

Suitability requirements

Resolution: NLT 3.5 between methocarbamol and guaifenesin, *System suitability solution*

Tailing factor: NMT 2.0, *Standard solution*

Relative standard deviation: NMT 5.0%, *Standard solution*

Analysis

Samples: *Standard solution* and *Sample solution*

Calculate the percentage of each impurity in the portion of Methocarbamol taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times (1/F) \times 100$$

r_U = peak response of each impurity from the *Sample solution*

r_S = peak response of methocarbamol from the *Standard solution*

C_S = concentration of [USP Methocarbamol RS](#) in the *Standard solution* (mg/mL)

C_U = concentration of Methocarbamol in the *Sample solution* (mg/mL)

F = relative response factor (see [Table 1](#))

Acceptance criteria: See [Table 1](#).

Table 1

Name	Relative Retention Time	Relative Response Factor	Acceptance Criteria, NMT (%)
Guaifenesin	0.84	1.2	0.15
Methocarbamol isomer ^a	0.90	1.0	0.05
Methocarbamol	1.0	—	—
Methocarbamol dioxolone ^b	1.3	1.0	0.05
Any individual unspecified impurity	—	—	0.05
Total impurities	—	—	1.0

^a 1-Hydroxy-3-(2-methoxyphenoxy)propan-2-yl carbamate.

^b 4-[(2-Methoxyphenoxy)methyl]-1,3-dioxolan-2-one.

SPECIFIC TESTS

- **LOSS ON DRYING (731).**

Analysis: Dry at 60° for 2 h.

Acceptance criteria: NMT 0.5%

ADDITIONAL REQUIREMENTS

- **PACKAGING AND STORAGE:** Preserve in tight containers. Store at controlled room temperature.
- **USP REFERENCE STANDARDS (11).**

Auxiliary Information - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
METHOCARBAMOL	Documentary Standards Support	SM42020 Small Molecules 4

Chromatographic Database Information: [Chromatographic Database](#)

Most Recently Appeared In:

Pharmacopeial Forum: Volume No. PF 41(4)

Current DocID: GUID-91439311-E1CE-4101-A135-52A0D5DC89E1_4_en-US

DOI: https://doi.org/10.31003/USPNF_M51200_04_01

DOI ref: [74000](#)

OFFICIAL