

Status: Currently Official on 15-Feb-2025
Official Date: Official as of 01-May-2020
Document Type: USP Monographs
DocId: GUID-53E05050-19AC-4BFD-B3CB-CFF800AAC222_6_en-US
DOI: https://doi.org/10.31003/USPNF_M51450_06_01
DOI Ref: 63jlk

© 2025 USPC
Do not distribute

Methotrexate Injection

To view the Notice from the Expert Committee that posted in conjunction with this accelerated revision, please click

<https://www.uspnf.com/rb-methotrexate-injection-20190501>

DEFINITION

Methotrexate Injection is a sterile solution of Methotrexate in Water for Injection prepared with the aid of Sodium Hydroxide. It contains NLT 90.0% and NMT 110.0% of the labeled amount of methotrexate ($C_{20}H_{22}N_8O_5$).

IDENTIFICATION

Change to read:

- A. **▲SPECTROSCOPIC IDENTIFICATION TESTS (197), Infrared Spectroscopy: (197K)▲** (CN 1-May-2020)

Sample: Dilute, if necessary, a volume of Injection, equivalent to about 25 mg of methotrexate, with water to obtain a solution with a concentration of about 2.5 mg/mL. Adjust with 0.1 N hydrochloric acid to a pH of 4.0. Place the slurry in a 50-mL centrifuge tube, and centrifuge. Decant the supernatant, add 25 mL of acetone, shake, and pass through a solvent-resistant membrane filter of 0.45- μ m pore size. Air-dry the filtered precipitate.

Acceptance criteria: Meets the requirements

- B. The retention time of the major peak of the *Sample solution* corresponds to that of the *Standard solution*, as obtained in the Assay.

ASSAY

- **PROCEDURE**

Buffer: 3.4 mg/mL of anhydrous monobasic sodium phosphate in water. Adjust with 1 N sodium hydroxide to a pH of 6.0.

Solution A: Acetonitrile and *Buffer* (5:95)

Solution B: Acetonitrile and *Buffer* (50:50)

Mobile phase: See [Table 1](#).

Table 1

Time (min)	Solution A (%)	Solution B (%)
0.0	100	0
30	50	50
34	50	50
35	100	0
40	100	0

Standard solution: 0.2 mg/mL of [USP Methotrexate RS](#) in *Solution A* prepared as follows. Add a sufficient amount of [USP Methotrexate RS](#) to a suitable volumetric flask and add dimethyl sulfoxide equivalent to 5% of the flask volume. Sonicate to achieve dissolution, then dilute with *Solution A* to volume.

Sample solution: Nominally 0.2 mg/mL of methotrexate from Injection prepared as follows. Transfer a sufficient amount of Injection to an appropriate volumetric flask. Add about 5% of the flask volume of dimethyl sulfoxide and sonicate for 2 min at ambient temperature, then add 30% of the flask volume of *Solution A* and sonicate. Dilute with *Solution A* to volume.

Chromatographic system

(See [Chromatography \(621\), System Suitability](#).)

Mode: LC

Detector: UV 280 nm

Column: 4.6-mm \times 25-cm; 5- μ m packing L1

Flow rate: 1 mL/min

Injection volume: 20 μ L

System suitability**Sample:** Standard solution**Suitability requirements****Tailing factor:** NMT 2.0**Relative standard deviation:** NMT 2.0%**Analysis****Samples:** Standard solution and Sample solutionCalculate the percentage of the labeled amount of methotrexate ($C_{20}H_{22}N_8O_5$) in the portion of Injection taken:

$$\text{Result} = (r_u/r_s) \times (C_s/C_u) \times 100$$

 r_u = peak response from the Sample solution r_s = peak response from the Standard solution C_s = concentration of [USP Methotrexate RS](#) in the Standard solution ($\mu\text{g/mL}$) C_u = nominal concentration of methotrexate in the Sample solution ($\mu\text{g/mL}$)**Acceptance criteria:** 90.0%–110.0%**IMPURITIES****Change to read:****• ORGANIC IMPURITIES****Solution A, Solution B, Mobile phase, Sample solution, and Chromatographic system:** Proceed as directed in the Assay.**Standard solution:** 0.2 $\mu\text{g/mL}$ each of [USP Methotrexate RS](#), [USP Methotrexate Related Compound B RS](#), [USP Methotrexate Related Compound C RS](#), and [USP Methotrexate Related Compound E RS](#) in Solution A prepared as follows. Add a sufficient amount of each Reference Standard to a suitable volumetric flask and add dimethyl sulfoxide equivalent to 5% of the flask volume. Sonicate to achieve dissolution, then dilute with Solution A to volume. Sonicate if necessary to aid dissolution.**Sample solution:** Nominally 0.2 mg/mL of methotrexate from Injection prepared as directed in the Assay**System suitability****Sample:** Standard solution[NOTE—See [Table 2](#) for relative retention times.]**Suitability requirements****Resolution:** NLT 1.5 between methotrexate related compound B and methotrexate related compound C**Relative standard deviation:** NMT 5.0% each for methotrexate, methotrexate related compound B, methotrexate related compound C, and methotrexate related compound E**Analysis****Samples:** Standard solution and Sample solution

Calculate the percentage of methotrexate related compound B and methotrexate related compound C in the portion of Injection taken:

$$\text{Result} = (r_u/r_s) \times (C_s/C_u) \times 100$$

 r_u = peak response of each corresponding impurity from the Sample solution r_s = peak response of each corresponding Reference Standard from the Standard solution C_s = concentration of each corresponding Reference Standard in the Standard solution ($\mu\text{g/mL}$) C_u = nominal concentration of methotrexate in the Sample solution ($\mu\text{g/mL}$)

Calculate the percentage of methotrexate related compound E free acid in the portion of Injection taken:

$$\text{Result} = (r_u/r_s) \times (C_s/C_u) \times (M_{r1}/M_{r2}) \times 100$$

 r_u = peak response from the Sample solution r_s = peak response from the Standard solution C_s = concentration of [USP Methotrexate Related Compound E RS](#) in the Standard solution ($\mu\text{g/mL}$) C_u = nominal concentration of methotrexate in the Sample solution ($\mu\text{g/mL}$) M_{r1} = molecular weight of methotrexate related compound E free acid, 325.33 M_{r2} = molecular weight of [USP Methotrexate Related Compound E RS](#), 343.56

[NOTE—[USP Methotrexate Related Compound E RS](#) is 4-[(2,4-Diaminopteridin-6-yl)methyl](methyl)amino}benzoic acid, hemihydrochloride.]

Calculate the percentage of any individual unspecified degradation product in the portion of Injection taken:

$$\text{Result} = (r_U/r_S) \times (C_S/C_U) \times 100$$

 r_U = peak response of each unspecified degradation product from the *Sample solution* r_S = peak response of methotrexate from the *Standard solution* C_S = concentration of [USP Methotrexate RS](#) in the *Standard solution* (μg/mL) C_U = nominal concentration of methotrexate in the *Sample solution* (μg/mL)**Acceptance criteria:** See [Table 2](#). The reporting threshold is 0.1%.**Table 2**

Name	Relative Retention Time	Acceptance Criteria, NMT (%)
Methotrexate related compound B	0.67	0.3
Methotrexate related compound C	0.73	▲4.0▲ (RB 1-May-2019)
Methotrexate	1.0	—
Methotrexate related compound E free acid ^a	1.41	0.3
Any individual unspecified degradation product	—	0.2
Total unspecified degradation products	—	1.0

^a 4-[(2,4-Diaminopteridin-6-yl)methyl](methyl)amino}benzoic acid.**SPECIFIC TESTS**

- [pH \(791\)](#): 7.0–9.0
- [BACTERIAL ENDOTOXINS TEST \(85\)](#): NMT 0.4 USP Endotoxin Units/mg of methotrexate sodium
- **OTHER REQUIREMENTS:** Meets the requirements in [Injections and Implanted Drug Products \(1\)](#).

ADDITIONAL REQUIREMENTS

- **PACKAGING AND STORAGE:** Preserve in single-dose or in multiple-dose containers, preferably of Type I glass, protected from light. Store at controlled room temperature.

- [USP REFERENCE STANDARDS \(11\)](#)

[USP Methotrexate RS](#)[USP Methotrexate Related Compound B RS](#)

(S)-2-(4-[(2,4-Diaminopteridin-6-yl)methylamino]benzamido)pentanedioic acid.

 $C_{19}H_{20}N_8O_5$ 440.41[USP Methotrexate Related Compound C RS](#)

(S)-2-(4-[(2-Amino-4-oxo-1,4-dihydropteridin-6-yl)methyl](methyl)amino}benzamido)pentanedioic acid.

 $C_{20}H_{21}N_7O_6$ 455.42[USP Methotrexate Related Compound E RS](#)

4-[(2,4-Diaminopteridin-6-yl)methyl](methyl)amino}benzoic acid, hemihydrochloride.

 $C_{15}H_{15}N_7O_2 \cdot \frac{1}{2}HCl$ 343.56 (anhydrous) $C_{15}H_{15}N_7O_2$ 325.33 (free acid)**Auxiliary Information** - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
METHOTREXATE INJECTION	Documentary Standards Support	SM32020 Small Molecules 3

Most Recently Appeared In:

Pharmacopeial Forum: Volume No. PF 41(4)

Current DocID: GUID-53E05050-19AC-4BFD-B3CB-CFF800AAC222_6_en-US

DOI: https://doi.org/10.31003/USPNF_M51450_06_01

DOI ref: [63jlk](#)

OFFICIAL