

Status: Currently Official on 16-Feb-2025  
Official Date: Official Prior to 2013  
Document Type: Reagents  
DocId: GUID-614FBAB8-29C5-4F56-B373-F42F2F542D3D\_1\_en-US  
DOI: [https://doi.org/10.31003/USPNF\\_R2215\\_01\\_01](https://doi.org/10.31003/USPNF_R2215_01_01)  
DOI Ref: sr80g

© 2025 USPC  
Do not distribute

## $\beta$ -Naphthoquinone-4-sodium Sulfonate,

$C_{10}H_5NaO_5S$  260.20—Yellow to orange-yellow crystals or crystalline powder. Soluble in about 10 parts of water; insoluble in alcohol.

**Loss on Drying (731):** Dry it in vacuum at about 50°: it loses not more than 2.0% of its weight.

**Residue on Ignition** (Reagent test): Ignite 1 g of dried sample with 3 mL of sulfuric acid: the residue weighs between 265 and 280 mg (between 26.5% and 28.0%).

**Auxiliary Information** - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
SS-NAPHTHOQUINONE-4-SODIUM SULFONATE	<a href="#">Margareth R.C. Marques</a> Principal Scientific Liaison	HDQ Headquarters

**Most Recently Appeared In:**

Pharmacopeial Forum: Volume No. Information currently unavailable

**Current DocId:** [GUID-614FBAB8-29C5-4F56-B373-F42F2F542D3D\\_1\\_en-US](#)

**DOI:** [https://doi.org/10.31003/USPNF\\_R2215\\_01\\_01](https://doi.org/10.31003/USPNF_R2215_01_01)

**DOI ref:** [sr80g](#)

