

Status: Currently Official on 16-Feb-2025

Official Date: Official as of 01-Aug-2017

Document Type: USP Monographs

DocId: GUID-BF5407E0-C675-4CFD-A3BB-21DB0C789397_1_en-US

DOI: https://doi.org/10.31003/USPNF_M56131_01_01

DOI Ref: e1mnz

© 2025 USPC

Do not distribute

Neomycin and Polymyxin B Sulfates, Bacitracin, and Hydrocortisone Acetate Ophthalmic Ointment

DEFINITION

Neomycin and Polymyxin B Sulfates, Bacitracin, and Hydrocortisone Acetate Ophthalmic Ointment contains the equivalent of NLT 90.0% and NMT 140.0% of the labeled amounts of neomycin, polymyxin B, and bacitracin, and NLT 90.0% and NMT 110.0% of the labeled amount of hydrocortisone acetate in a suitable ointment base.

IDENTIFICATION

- **A. THIN-LAYER CHROMATOGRAPHIC IDENTIFICATION TEST (201BNP):** Meets the requirements
- **B.** The retention time of the hydrocortisone acetate peak of the *Sample solution* corresponds to that of the *Standard solution*, as obtained in the Assay for *Hydrocortisone Acetate*.

ASSAY

• NEOMYCIN

(See [Antibiotics—Microbial Assays \(81\)](#).)

Sample solution: Shake a portion of Ophthalmic Ointment in a separator with 50 mL of [ether](#). Extract with four 20-mL portions of *Buffer B.3*. Combine the aqueous extracts, and dilute with *Buffer B.3* to a suitable volume.

Analysis: Proceed as directed in the chapter. Dilute the *Sample solution* with *Buffer B.3* to obtain a *Test Dilution* having a neomycin concentration that is nominally equivalent to the median level of the standard.

Acceptance criteria: 90.0%–140.0%

• POLYMYXIN B

(See [Antibiotics—Microbial Assays \(81\)](#).)

Sample solution: Shake a portion of Ophthalmic Ointment with 50 mL of [ether](#) in a separator. Extract with four 25-mL portions of *Buffer B.6*. Combine the aqueous extracts, and dilute with *Buffer B.6* to a suitable volume.

Analysis: Proceed as directed in the chapter. Dilute the *Sample solution* with *Buffer B.6* to obtain a *Test Dilution* having a concentration that is nominally equivalent to the median level of the standard (10 polymyxin B units/mL). Add to each *Test Dilution* of the standard a quantity of [USP Neomycin Sulfate RS](#), dissolved in *Buffer B.6*, to obtain the same concentration of neomycin as in the *Test Dilution* of the sample.

Acceptance criteria: 90.0%–140.0%

• BACITRACIN

(See [Antibiotics—Microbial Assays \(81\)](#).)

Sample solution: Shake a portion of Ophthalmic Ointment with 50 mL of [ether](#) in a separator. Extract with four 20-mL portions of *Buffer B.1*. Combine the buffer extracts, and dilute with *Buffer B.1* to a suitable volume.

Analysis: Proceed as directed in the chapter. Add sufficient [0.01 N hydrochloric acid](#) to a portion of the *Sample solution* so that the amount of hydrochloric acid in the *Test Dilution* is the same as in the median level of the standard. Dilute with *Buffer B.1* to obtain a *Test Dilution* having a bacitracin concentration that is nominally equivalent to the median level of the standard.

Acceptance criteria: 90.0%–140.0%

• HYDROCORTISONE ACETATE

Mobile phase: [Butyl chloride](#), water-saturated [butyl chloride](#), [tetrahydrofuran](#), [methanol](#), and [glacial acetic acid](#) (475:475:70:35:30)

Standard solution: 0.10 mg/mL of [USP Hydrocortisone Acetate RS](#) in water-saturated [chloroform](#)

Sample solution: Nominally 0.10 mg/mL of hydrocortisone acetate from Ophthalmic Ointment prepared as follows. Transfer a portion of Ophthalmic Ointment containing nominally 2.5 mg of hydrocortisone acetate to a closable container. Add 25.0 mL of water-saturated [chloroform](#) and about 10 glass beads. Securely close the container, and shake vigorously for approximately 15 min. Centrifuge, and use the clear, lower chloroform layer.

Chromatographic system

(See [Chromatography \(621\), System Suitability](#).)

Mode: LC

Detector: UV 254 nm**Column:** 3.9-mm × 30-cm; 10-µm packing [L3](#)**System suitability****Sample:** Standard solution**Suitability requirements****Relative standard deviation:** NMT 2.0%**Analysis****Samples:** Standard solution and Sample solutionCalculate the percentage of the labeled amount of hydrocortisone acetate ($C_{23}H_{32}O_6$) in the portion of Ophthalmic Ointment taken:

$$\text{Result} = (r_u/r_s) \times (C_s/C_u) \times 100$$

 r_u = peak response from the Sample solution r_s = peak response from the Standard solution C_s = concentration of [USP Hydrocortisone Acetate RS](#) in the Standard solution (mg/mL) C_u = nominal concentration of hydrocortisone acetate in the Sample solution (mg/mL)**Acceptance criteria:** 90.0%–110.0%**SPECIFIC TESTS**

- [STERILITY TESTS \(71\)](#): Meets the requirements
- **OTHER REQUIREMENTS:** It meets the requirements for Particulate and Foreign Matter and Container Contents in [Ophthalmic Products—Quality Tests \(771\)](#), [Drug Product Quality, Universal Tests, Particulate and Foreign Matter](#) and [Container Contents](#).

ADDITIONAL REQUIREMENTS

- **PACKAGING AND STORAGE:** Preserve in collapsible ophthalmic ointment tubes.

- [USP REFERENCE STANDARDS \(11\)](#):

[USP Bacitracin Zinc RS](#)[USP Hydrocortisone Acetate RS](#)[USP Neomycin Sulfate RS](#)[USP Polymyxin B Sulfate RS](#)**Auxiliary Information** - Please [check for your question in the FAQs](#) before contacting USP.

Topic/Question	Contact	Expert Committee
NEOMYCIN AND POLYMYXIN B SULFATES, BACITRACIN, AND HYDROCORTISONE ACETATE OPHTHALMIC OINTMENT	Julie Zhang Associate Science & Standards Liaison	BIO42020 Biologics Monographs 4 - Antibiotics
REFERENCE STANDARD SUPPORT	RS Technical Services RSTECH@usp.org	BIO42020 Biologics Monographs 4 - Antibiotics

Chromatographic Database Information: [Chromatographic Database](#)**Most Recently Appeared In:**

Pharmacopeial Forum: Volume No. PF 39(5)

Current DocID: [GUID-BF5407E0-C675-4CFD-A3BB-21DB0C789397_1_en-US](#)**DOI:** https://doi.org/10.31003/USPNF_M56131_01_01**DOI ref:** [e1mnz](#)